

**Lecture notes for the introductory talk on the single-reference many-body perturbation theory prepared by Piotr Piecuch for the Workshop of the *Espace de Structure et de Réactions Nucléaires Théorique* on “Many-Body Perturbation Theories in Modern Quantum Chemistry and Nuclear Physics,” March 26-30, 2018, CEA Saclay, Gif-sur-Yvette, France.**

-+-

# MANY-BODY PERTURBATION THEORY (SINGLE-REFERENCE CASE).

## 1. Introductory remarks.

We will use the Rayleigh-Schrödinger perturbation theory (RSPD) for a non-degenerate ground state to solve the many-particle (many-Fermion) Schrödinger equation,

$$H|\Psi\rangle = E_0|\Psi\rangle, \quad (1)$$

where the correlated ground state  $|\Psi\rangle$  can be obtained by perturbing the independent-particle-model (IPM) single-determinantal state  $|\tilde{\Psi}\rangle$  that will also serve as a Fermi vacuum. We will assume that our Hamiltonian  $H$  consists of one- and two-body components,  $Z$  and  $V$ , respectively, so that

$$H = Z + V, \quad (2)$$

where

$$Z = \sum_{p,q} \langle p | \hat{z} | q \rangle X_p^+ X_q \quad (3)$$

and

$$V = \frac{1}{2} \sum_{pq,rs} \langle pq|\hat{v}|rs\rangle X_p^+ X_q^+ X_s^- X_r^- \quad (4)$$

$$= \frac{1}{4} \sum_{pq,rs} \langle pq|\hat{v}|rs\rangle A \quad X_p^+ X_q^+ X_s^- X_r^-,$$

with

$$\langle pq|\hat{v}|rs\rangle_A = \langle pq|\hat{v}|rs\rangle - \langle pq|\hat{v}|sr\rangle \quad (5)$$

representing the antisymmetrized matrix elements.

Here,  $X_p^+$  ( $X_p$ ) are the creation (annihilation) operators associated with single-particle states (in quantum chemistry, spin-orbitals)  $|p\rangle$ .

To facilitate our considerations, where it will be assumed that  $|\Psi_0\rangle$  is obtained by perturbing the 1PM Fermi vacuum state  $|\Psi_0\rangle$  ( $|\Psi_0\rangle$  could, for example, be a Hartree-Fock determinant, although other choices are certainly possible), we will focus on the Schrödinger equation written as

$$H_N |\Psi_0\rangle = \Delta E_0 |\Psi_0\rangle, \quad (6)$$

where

$$H_N = H - \langle \Psi_0 | H | \Psi_0 \rangle \quad (7)$$

-3-

is the Hamiltonian in the normal-ordered form  
(we will return to this later) and

$$\Delta E_0 = E_0 - \langle \Psi_0 | H(\Psi_0) \rangle. \quad (8)$$

If  $|\Psi_0\rangle$  is a Hartree-Fock state,  $\Delta E_0$  is the conventional correlation energy. We will be seeking the solutions of Eq. (8), where we know that the exact  $|\Psi\rangle$  can be written as

$$\begin{aligned} |\Psi\rangle &= |\Psi_0\rangle + \sum_{i,a} c_a^i |\Psi_i^a\rangle + \\ &+ \sum_{i < j, a < b} c_{ab}^{ij} |\Psi_{ij}^{ab}\rangle + \dots \\ &= |\Psi_0\rangle + \sum_{n=1}^N \sum_{\substack{i_1 < \dots < i_n \\ a_1 < \dots < a_n}} c_{a_1 \dots a_n}^{i_1 \dots i_n} |\Psi_{i_1 \dots i_n}^{a_1 \dots a_n}\rangle \end{aligned} \quad (9)$$

where

$$|\Psi_i^a\rangle = X_a^\dagger X_i |\Psi_0\rangle \equiv E_i^a |\Psi_0\rangle,$$

$$|\Psi_{ij}^{ab}\rangle = X_a^\dagger X_i X_b^\dagger X_j |\Psi_0\rangle = E_{ij}^{ab} |\Psi_0\rangle,$$

$$\begin{aligned} |\Psi_{i_1 \dots i_n}^{a_1 \dots a_n}\rangle &= \prod_{g=1}^n X_{a_g}^\dagger X_{i_g} |\Psi_0\rangle \\ &= \prod_{g=1}^n E_{i_g}^{a_g} |\Psi_0\rangle = E_{i_1 \dots i_n}^{a_1 \dots a_n} |\Psi_0\rangle \end{aligned} \quad (10)$$

one-particle-one-hole

are the  $1p-1h$ ,  $2p-2h$ , ...,  $n p-nh$  excited determinants, in the form of a perturbative expansion,

$$|\bar{\Psi}_0\rangle = |\bar{\Psi}_0^{(0)}\rangle + |\bar{\Psi}_0^{(1)}\rangle + |\bar{\Psi}_0^{(2)}\rangle + \dots$$

$$= \sum_{n=0}^{\infty} |\bar{\Psi}_0^{(n)}\rangle, \quad (11)$$

in which  $|\bar{\Psi}_0^{(0)}\rangle = |\bar{\Psi}_0\rangle$  and  $|\bar{\Psi}_0^{(n)}\rangle$  with  $n > 1$  are the corrections to the zeroth-order state  $|\bar{\Psi}_0\rangle$ . The corresponding correlation energy (defined as  $E_0$  minus  $\langle \bar{\Psi}_0 | \hat{H} | \bar{\Psi}_0 \rangle$ ) will be represented as

$$\Delta E_0 = \Delta E_0^{(0)} + \Delta E_0^{(1)} + \Delta E_0^{(2)} + \dots$$

$$= \sum_{n=0}^{\infty} \Delta E_0^{(n)}, \quad (12)$$

where, quite obviously and as we will see,  $\Delta E_0^{(0)} = 0$  and  $\Delta E_0^{(1)} = 0$ . We will use the RSPT approach to determine expansions (11) and (12). Before doing this, let us discuss key elements of RSPT for a generic Hermitian eigenvalue problem,

$$K|\bar{\Psi}\rangle = k_0|\bar{\Psi}\rangle \quad (13)$$

for a non-degenerate state  $|\bar{\Psi}_0\rangle$ .

2. Rayleigh-Schrödinger perturbation theory for a nondegenerate eigenvalue problem.

We want to solve

$$K|\Psi\rangle = k_0|\Psi\rangle. \quad (14)$$

In RSPT, we assume that we can split  $K$  into the unperturbed part  $K_0$  and perturbation  $W$ ,

$$K = K_0 + W, \quad (15)$$

such that we know all eigenvalues  $\varepsilon_n$  and all eigenstates  $|\Phi_n\rangle$  of  $K_0$ ,

$$K_0|\Phi_n\rangle = \varepsilon_n|\Phi_n\rangle, \quad n=0,1,2,\dots \quad (16)$$

$K_0$  is Hermitian, so states  $|\Phi_n\rangle$  form an orthonormal basis in the Hilbert space,

$$\langle \Phi_m | \Phi_n \rangle = \delta_{mn}, \quad (17)$$

and  $\varepsilon_n$ 's are real numbers. We seek the solution of Eq. (14) in the form

$$|\Psi\rangle = \mathcal{S}|\Phi\rangle,$$

where  $\mathcal{S}$  is the so-called wave operator,

-6-

using intermediate normalization,

$$\langle \psi_0 | \bar{\psi}_0 \rangle = \langle \psi_0 | S_2 | \bar{\psi}_0 \rangle = \langle \bar{\psi}_0 | \bar{\psi}_0 \rangle = 1. \quad (18)$$

We obtain,  
projecting both sides  
on  $\langle \psi_0 |$

$$\langle \psi_0 | (K_0 + W) | \bar{\psi}_0 \rangle = k_0 \langle \bar{\psi}_0 | \bar{\psi}_0 \rangle,$$
$$\underbrace{\langle \psi_0 | K_0 | \bar{\psi}_0 \rangle}_{\propto \langle \psi_0 |} + \langle \psi_0 | W | \bar{\psi}_0 \rangle = k_0 \langle \bar{\psi}_0 | \bar{\psi}_0 \rangle = k_0,$$

$$k_0 = \propto_0 + \langle \psi_0 | W | \bar{\psi}_0 \rangle$$
$$= \propto_0 + \langle \psi_0 | W S_2 | \bar{\psi}_0 \rangle$$
$$= \propto_0 + \langle \psi_0 | \bar{\tau} | \bar{\psi}_0 \rangle, \quad (19)$$

where

$$\bar{\tau} = PWS_2,$$

with

$$P = |\bar{\psi}_0 \rangle \langle \bar{\psi}_0| \quad (20)$$

representing the projection operator on the P-space spanned by  $|\bar{\psi}_0 \rangle$ , is the so-called reaction operator.

-7-

$S_2$  maps  $\mathcal{H}_0$  onto  $\mathcal{H}_0$ , but without knowing how it acts on the remaining basis states  $\{|E_n\rangle\}$ , with  $n \geq 1$ , it is not uniquely defined. RSPT is one of the infinitely many possibilities of finding  $S_2$ . The key quantity for setting up the RSPT series is the REDUCED RESOLVENT.

To define the reduced resolvent, we decompose the Hilbert space  $\mathcal{H}$  into the space spanned by  $\mathcal{H}_0$  and the orthogonal complement called the  $\mathcal{H}_Q$  space, so that

$$\mathcal{H} = \mathcal{H}_0 \oplus \mathcal{H}_Q. \quad (21)$$

The corresponding projection operators are  $P$ , Eq. (20), and

$$Q = I - P = \sum_{n=1}^{\infty} |\psi_n\rangle \langle \psi_n|. \quad (22)$$

The reduced resolvent of operator  $K_0$  which is parameterized by  $\alpha$ , is formally defined as

[ $\alpha$ -real variable]

$$R_\alpha(K_0) = Q[\alpha P + Q(\alpha - K_0)Q]^{-1}Q,$$

where  $\alpha \neq 0$ . It is easy to show that the matrix representation of  $R_\alpha(K_0)$  in a basis set (23)

-8-

defined by  $\{\mathbb{E}_n\}$ ,  $n=0, 1, 2, \dots$ , is

$P$ -space  $\rightarrow \begin{pmatrix} 0 & | & 0 \\ \hline - & - & - \end{pmatrix} \equiv$

$Q$ -space  $\rightarrow \begin{pmatrix} 0 & | & (xI_Q - QKQ)^{-1} \\ \hline \equiv & & \end{pmatrix}.$  (24)

$\uparrow_{P\text{-space}} \quad \uparrow_{Q\text{-space}}$

The spectral representation of  $R_{\alpha}(K)$  is

$$R_{\alpha}(K) = \sum_{n=1}^{\infty} \frac{[\mathbb{E}_n \times \mathbb{E}_n]}{\alpha - \alpha_n}. \quad (25)$$

eigenvalues of  $QKQ$

Thus,  $R_{\alpha}(K)$  becomes singular for  $\alpha = \alpha_1, \alpha_2, \dots$ ,  
but not for  $\alpha = \alpha_0$ , since  $\alpha_0$  is non-degenerate (by  
assumption).

Properties of  $P$ ,  $Q$ , and  $R_{\alpha}(K)$ :

- $P^2 = P$ ,  $P^t = P$ ,  $Q^2 = Q$ ,  $Q^t = Q$ ,  $PQ = QP = 0$ ,  
(idempotent)      (hermitian)
- $R_{\alpha}(K) = R_{\alpha}(K)^t$  (for real  $\alpha$ ),
- $R_{\alpha}(K)P = PR_{\alpha}(K) = 0$ .
- $Q R_{\alpha}(K) = R_{\alpha}(K)Q = R_{\alpha}(K)$ .

-3-

this is why sometimes we  
 write  $R_x(K) = \frac{Q}{x-K_0}$

$$\bullet Q(x-K_0) R_x(K_0) = R_x(K_0)(x-K_0)$$

$$= Q. \quad (26)$$

There are other. The last one is particularly important, and we can prove it as follows:

$$Q(x-K_0) R_x(K_0) = Q \sum_{n=1}^{\infty} \frac{(x-K_0) |z_n \rangle \langle z_n|}{x-z_n}$$

$$= Q \sum_{n=1}^{\infty} \frac{(x-x_n) |z_n \rangle \langle z_n|}{x-x_n} = Q \cdot Q = Q. \quad (27)$$

Equipped with the above definitions, we define the reduced resolvent of  $K_0$  at  $x=x_0$ , which I will call the upperended reduced resolvent,

$$R^{(0)} = R_{x=x_0}(K_0) = Q [P + Q(x_0 - K_0) P] Q$$

$$= \sum_{n=1}^{\infty} \frac{|z_n^{(0)}\rangle \langle z_n^{(0)}|}{x_0 - z_n}. \quad (28)$$

We can use it to develop the RSPT series in the following few steps:

(i) We know that

$$|z\rangle = (P+Q)|z\rangle = |z\rangle \otimes |z\rangle$$

$$+ Q|z\rangle = |z\rangle + Q|z\rangle. \quad (29)$$

-10

We consider the following expression:

$$\begin{aligned}
 (\chi_0 - K_0) \underline{Q} |\Psi\rangle &= Q(\chi_0 - K_0) |\Psi\rangle \\
 &= Q(\chi_0 - K + W) |\Psi\rangle \\
 &= Q(\chi_0 - k_0 + W) |\Psi\rangle, \tag{30}
 \end{aligned}$$

where we used the fact that  $[Q, K_0] = 0$  (obvious). Let us define

$$W' = W - (k_0 - \chi_0). \tag{31}$$

We obtain,

$$(\chi_0 - K_0) \underline{Q} |\Psi\rangle = Q W' |\Psi\rangle. \tag{32}$$

(ii) We know that (see Eq. (26))

$$\begin{aligned}
 Q(\chi_0 - K_0) R^{(0)} &= R^{(0)}(\chi_0 - K_0) Q \\
 &= Q. \tag{33}
 \end{aligned}$$

Thus, from Eqs. (32) and (33), we obtain,

$$\underbrace{R^{(0)}(\chi_0 - K_0) Q}_{\text{Eq.(33)} \rightarrow Q} |\Psi\rangle = \underbrace{R^{(0)} Q W'}_{\text{see p.8} \rightarrow R^{(0)}} |\Psi\rangle, \tag{34}$$

- 11 -

$$Q|\psi_0\rangle = R^{(0)} W' |\psi_0\rangle, \quad (35)$$

$$\begin{aligned} |\psi\rangle &= |\psi_0\rangle + Q|\psi_0\rangle \\ &= |\psi_0\rangle + R^{(0)} W' |\psi_0\rangle. \end{aligned} \quad (36)$$

(iii) Hermiting the last relationship, we obtain,

$$\begin{aligned} |\psi\rangle &= |\psi_0\rangle + R^{(0)} W' (|\psi_0\rangle + R^{(0)} W' |\psi_0\rangle) \\ &= |\psi_0\rangle + (R^{(0)} W') |\psi_0\rangle + (R^{(0)} W')^2 |\psi_0\rangle \\ &= \dots = \sum_{n=0}^{\infty} (R^{(0)} W')^n |\psi_0\rangle. \end{aligned} \quad (37)$$

Thus,  $|\psi\rangle = \sum_{n=0}^{\infty} (R^{(0)} W')^n |\psi_0\rangle, \quad (38)$

where  $W'$  is given by Eq. (31).

Using Eq. (19), we obtain

$$\begin{aligned} k_0 &= \alpha_0 + \langle \psi_0 | W |\psi\rangle = \alpha_0 \\ &+ \sum_{n=0}^{\infty} \langle \psi_0 | W (R^{(0)} W')^n |\psi_0\rangle. \end{aligned} \quad (39)$$

We can make further slight simplifications,

$$\begin{aligned}
 |\bar{\psi}_0\rangle &= \sum_{n=0}^{\infty} (R^{(0)} W')^n |\bar{\psi}\rangle \\
 &= |\bar{\psi}_0\rangle + \sum_{n=1}^{\infty} (R^{(0)} W')^n |\bar{\psi}\rangle \\
 &= |\bar{\psi}_0\rangle + \sum_{n=1}^{\infty} (R^{(0)} W')^n (R^{(0)} W') |\bar{\psi}\rangle \\
 &= |\bar{\psi}_0\rangle + \sum_{n=0}^{n=0} (R^{(0)} W')^n (R^{(0)} W) |\bar{\psi}\rangle,
 \end{aligned} \tag{40}$$

since

$$\begin{aligned}
 R^{(0)} W' |\bar{\psi}\rangle &= R^{(0)} W |\bar{\psi}\rangle + R^{(0)} (\bar{\psi}_0 - k_0) |\bar{\psi}\rangle \\
 &= R^{(0)} W |\bar{\psi}\rangle
 \end{aligned} \tag{41}$$

$$(R^{(0)} |\bar{\psi}\rangle = R^{(0)} P(\bar{\psi}) = 0).$$

Similarly, and using Eqs. (19) and (40),

$$\begin{aligned}
 k_0 &= \bar{\psi}_0 + \langle \bar{\psi} | W |\bar{\psi}\rangle = \bar{\psi}_0 + \langle \bar{\psi}_0 | W |\bar{\psi}\rangle \\
 &\quad + \sum_{n=0}^{\infty} \langle \bar{\psi}_0 | W (R^{(0)} W')^n R^{(0)} W |\bar{\psi}\rangle.
 \end{aligned} \tag{42}$$

Summary:

$$|\tilde{\psi}_0\rangle = |\psi_0\rangle + \sum_{n=0}^{\infty} (R^{(0)}W')^n R^{(0)}W |\psi_0\rangle \quad (43a)$$

$$k_0 = \varepsilon_0 + \langle \tilde{\psi}_0 | W(\tilde{\psi}_0) \rangle + \sum_{n=0}^{\infty} \langle \tilde{\psi}_0 | W(R^{(0)}W')^n R^{(0)}W | \tilde{\psi}_0 \rangle \quad (43b)$$

or, using the wave and reaction operators,

$$|\tilde{\psi}_0\rangle = S\tilde{\psi}_0, \quad (44)$$

where

$$S\tilde{\psi} = P + \sum_{n=0}^{\infty} (R^{(0)}W')^n R^{(0)}WP, \quad (45)$$

and

$$k_0 = \varepsilon_0 + \langle \tilde{\psi}_0 | \tilde{\tau} | \tilde{\psi}_0 \rangle, \quad (46)$$

where

$$\tilde{\tau} = PW\tilde{S}\tilde{\psi} = PW\tilde{P} + \sum_{n=0}^{\infty} P(R^{(0)}W')^n R^{(0)}WP, \quad (47)$$

with

$$W' = W - (k_0 - \varepsilon_0).$$

$S^2$  used in the above equations is an example of Block wave operator, which satisfies

$$S^2 P = S^2, \quad PS^2 = P, \quad S^2 = S^2.$$

(or  $S^2 Q = S^2(1-P) = 0$ )

$\uparrow$   
 $S^2$  is idempotent (48)

The property  $PS^2 = P$  is an intermediate normalization condition, since

$$\langle \psi | \psi \rangle = \langle \psi | S^2 \psi \rangle = \langle \psi | PS^2 \psi \rangle$$

(48)


$$= \langle \psi | P \psi \rangle = \langle \psi | \psi \rangle = 1. \quad (49)$$

Because of  $S^2 = S^2$ ,  $S^2$  is sometimes called the non-orthogonal projector and we obtain this property as

$$S^2 = (S^2 P)(S^2 P) = S^2 (P S^2) P = S^2 P^2 = S^2 P$$

$$= S^2 \text{ (non-orthogonal, since } S^2 \neq S^2)^T \text{).} \quad (50)$$

model  
or  
reference  
space



target space

$$S^2 |\psi\rangle = |\psi\rangle$$

$$P |\psi\rangle = |\psi\rangle$$

$$S^2 |b\rangle = (P|b\rangle)^{-1}$$

$$S^2 b_1 = 0$$

(S^2 = 0)

$$M = \text{Span}\{|\psi\rangle\} \quad M = \text{Span}\{|\psi\rangle\}$$

When  $M_0$  is multi-dimensional, we obtain multi-reference theories, such as MR MBPT. In that case,  $M = \text{Span}\{|\psi\rangle\}_{j=1}^M$

-15-

In RSPT we define PT orders according to powers of  $W$  (zero-order:  $W^0$ , 1st order:  $W^1$ , second order:  $W^2$ , etc.). As a result,

$$\langle \overline{H_0} \rangle = \sum_{n=0}^{\infty} \langle \overline{H}_0^{(n)} \rangle, \\ k_0 = \sum_{n=0}^{\infty} k_0^{(n)}, \\ S^L = \sum_{n=0}^{\infty} S^L_n, \\ \tau = \sum_{n=0}^{\infty} \tau^{(n)} = \sum_{n=1}^{\infty} \tau^{(n)}, \text{ since} \\ \tau^{(n+1)} = PW S^L_n (\tau^{(0)} = 0). \quad (51)$$

In generating the above expansions, we must keep in mind that

$[W^1 \text{ contains 1st and higher-order terms}]$

$$W^1 = W - (k_0 - \delta_0) = W - \sum_{m=1}^{\infty} k_0^{(m)}, \\ \text{since } k_0^{(0)} = \delta_0. \quad [THIS IS WHY WE HAVE] \\ [RENORMALIZATION TERMS] \quad (52)$$

Using Eqs. (43) - (47) and Eq. (52), we obtain:

0-th order:  $\langle \overline{H}_0^{(0)} \rangle = \langle \overline{H}_0 \rangle \quad (S^L = P),$  (53)

as anticipated  $\rightarrow k_0^{(0)} = \delta_0 \quad (\tau^{(0)} = 0).$

-16-

1st order:  $|\tilde{\Psi}_0^{(1)}\rangle = R^{(0)}W|\tilde{\psi}_0\rangle$  ( $S^{(1)} = R^{(0)}WP$ ),  
 $k_0^{(1)} = \langle \tilde{\psi}_0 | W | \tilde{\psi}_0 \rangle$  ( $C^{(1)} = PW P$ ). (54)

2nd order: from  $n=1$  in (43a)

$$\begin{aligned} |\tilde{\Psi}_0^{(2)}\rangle &= R^{(0)}(W - k_0^{(1)})R^{(0)}W|\tilde{\psi}_0\rangle \\ &= (R^{(0)}W)^2|\tilde{\psi}_0\rangle - k_0^{(1)}R^{(0)2}W|\tilde{\psi}_0\rangle \quad (55) \\ &= (R^{(0)}W)^2|\tilde{\psi}_0\rangle - \langle \tilde{\psi}_0 | W | \tilde{\psi}_0 \rangle R^{(0)2}W|\tilde{\psi}_0\rangle, \end{aligned}$$

$$k_0^{(2)} = \langle \tilde{\psi}_0 | WR^{(0)}W|\tilde{\psi}_0\rangle. \quad \text{(from } n=0 \text{ in (43b)})$$

3rd order: from  $n=2$  in (43a)

$$\begin{aligned} |\tilde{\Psi}_0^{(3)}\rangle &= [R^{(0)}(W - k_0^{(1)})]^2 R^{(0)}W|\tilde{\psi}_0\rangle \\ &\quad - k_0^{(2)} R^{(0)2}W|\tilde{\psi}_0\rangle \quad \text{(from } n=1 \text{ in (43a))} \\ &= (R^{(0)}W)^3|\tilde{\psi}_0\rangle - \langle \tilde{\psi}_0 | W | \tilde{\psi}_0 \rangle \\ &\quad \times (R^{(0)2}WR^{(0)}W|\tilde{\psi}_0\rangle + R^{(0)}WR^{(0)2}W|\tilde{\psi}_0\rangle) \\ &\quad + \langle \tilde{\psi}_0 | W | \tilde{\psi}_0 \rangle^2 R^{(0)3}W|\tilde{\psi}_0\rangle \\ &\quad - \langle \tilde{\psi}_0 | WR^{(0)}W|\tilde{\psi}_0 \rangle R^{(0)2}W|\tilde{\psi}_0\rangle, \quad (56) \end{aligned}$$

$k_0^{(2)}$  from  $W'$

-17 - from  $n=1$  in (43b)

$$\begin{aligned}
 k_0^{(3)} &= \langle \psi_0 | W R^{(0)} (W - k_0^{(1)}) R^{(0)} W | \psi_0 \rangle \\
 &= \langle \psi_0 | W (R^{(0)} W)^2 | \psi_0 \rangle \\
 &\quad - \langle \psi_0 | W | \psi_0 \rangle \langle \psi_0 | W R^{(0)2} W | \psi_0 \rangle,
 \end{aligned}$$

etc. We should note that

$$\begin{aligned}
 k_0^{(n+1)} &= \langle \psi_0 | \tilde{\tau}^{(n+1)} | \psi_0 \rangle \\
 &= \langle \psi_0 | P W S^{(n)} | \psi_0 \rangle \\
 &= \langle \psi_0 | W | \tilde{\psi}_0^{(n)} \rangle, \quad (57)
 \end{aligned}$$

since  $\tilde{\tau}^{(n)} | \tilde{\psi}_0^{(n)} \rangle = | \tilde{\psi}_0^{(n)} \rangle$ . Thus we obtain  $k_0^{(n+1)}$  by attaching  $\langle \psi_0 | W$  to  $| \tilde{\psi}_0^{(n)} \rangle$ .

For example,

$$\begin{aligned}
 k_0^{(4)} &= \langle \psi_0 | W | \tilde{\psi}_0^{(3)} \rangle = \langle \psi_0 | W (R^{(0)} W)^3 | \psi_0 \rangle \\
 &\quad - \langle \psi_0 | W | \psi_0 \rangle \langle \psi_0 | W R^{(0)2} W R^{(0)} W | \psi_0 \rangle \\
 &\quad + \langle \psi_0 | W R^{(0)} W R^{(0)2} W | \psi_0 \rangle \\
 &\quad + \langle \psi_0 | W | \psi_0 \rangle^2 \langle \psi_0 | W R^{(0)3} W | \psi_0 \rangle \\
 &\quad - \langle \psi_0 | W R^{(0)} W | \psi_0 \rangle \langle \psi_0 | W R^{(0)2} W | \psi_0 \rangle. \quad (58)
 \end{aligned}$$

It is easy to show that the above equations combined with the spectral representation of  $R^{(0)}$  give the well-known RSP<sup>T</sup> corrections, for example,

$$\begin{aligned} \langle \tilde{\psi}^{(1)} \rangle &= \sum_{n=1}^{\infty} \frac{\langle \tilde{\psi}_0 | \tilde{\psi}_n \rangle \langle \tilde{\psi}_n | }{\tilde{\epsilon}_0 - \tilde{\epsilon}_n} W(\tilde{\psi}_0) \\ &= \sum_{n=1}^{\infty} \frac{\langle \tilde{\psi}_0 | W(\tilde{\psi}_n) }{\tilde{\epsilon}_0 - \tilde{\epsilon}_n} \langle \tilde{\psi}_n | , \end{aligned} \quad (59)$$

$$\begin{aligned} k_0^{(2)} &= \langle \tilde{\psi}_0 | W \sum_{n=1}^{\infty} \frac{\langle \tilde{\psi}_n | \tilde{\psi}_n \rangle}{\tilde{\epsilon}_0 - \tilde{\epsilon}_n} W(\tilde{\psi}_0) \rangle \\ &= \sum_{n=1}^{\infty} \frac{\langle \tilde{\psi}_0 | W(\tilde{\psi}_n) \langle \tilde{\psi}_n | W(\tilde{\psi}_0) \rangle}{\tilde{\epsilon}_0 - \tilde{\epsilon}_n}, \text{ etc.} \end{aligned} \quad (60)$$

In general,

$$\langle \tilde{\psi}^{(n)} \rangle = \underbrace{(R^{(0)}W)^n(\tilde{\psi}_0)}_{\text{principal term}} + \text{renormalization terms} \quad (61)$$

$$k_0^{(n+1)} = \underbrace{\langle \tilde{\psi}_0 | W(R^{(0)}W)^n(\tilde{\psi}_0)}_{+ \text{renormalization terms.}}$$

One can generate the renormalization terms using the Breit-Wigner-Huby Bracketing technique.

The idea is to insert non-stroddling bracket pairs  $\langle \rangle^{(0)}$  representing  $\langle \mathcal{S} | \dots | \mathcal{S} \rangle$  into the principal term and closing it such that the following rules are satisfied:

- no two brackets can touch
- bracketing operation including the rightmost end, in the case of eigenvalue corrections, the leftmost  $W$  are not allowed,
- each bracket must have  $W$  on each side (as in  $\langle W \dots W \rangle$ )

We assign the sign  $(-1)^r$  to a term with  $r$  inserted bracket pairs.

Examples:

$$\bullet k_0^{(3)} = \langle W (R^{(0)} W)^2 \rangle + \text{renorm. terms}$$

Principal term:

$$\langle W R^{(0)} W R^{(0)} W \rangle.$$

Renormalization terms (term, only one here):

$$\langle W R^{(0)} \cancel{\langle W \rangle} R^{(0)} W \rangle$$

$$= \langle W \rangle \langle W R^{(0)2} W \rangle. \text{ Sign } (-1)^1 = -1.$$

$$k_0^{(3)} = \langle W (R^{(0)} W)^2 \rangle - \langle W \rangle \langle W R^{(0)2} W \rangle. (\pi=1)$$

-20-

•  $K_0^{(4)} = \langle W (R^{\alpha} W)^3 \rangle + \text{renorm. terms.}$

Principal term :

$$\langle W R^{\alpha} W R^{\alpha} W R^{\alpha} W \rangle.$$

Renormalization terms :

$$\langle W R^{\alpha} W R^{\alpha} W R^{\alpha} W \rangle \quad (\alpha=1)$$

$$\langle W R^{\alpha} W R^{\alpha} W R^{\alpha} W \rangle \quad (\alpha=1)$$

$$\langle W R^{\alpha} W R^{\alpha} W R^{\alpha} W R^{\alpha} W \rangle \quad (\alpha=2)$$

$$\langle W R^{\alpha} W R^{\alpha} W R^{\alpha} W R^{\alpha} W \rangle \quad (\alpha=1)$$

$$K_0^{(4)} = \langle W (R^{\alpha} W)^3 \rangle$$

$$- \langle W \rangle \langle W R^{\alpha 2} W R^{\alpha} W \rangle$$

$$- \langle W \rangle \langle W R^{\alpha} W R^{\alpha 2} W \rangle$$

$$+ \langle W \rangle^2 \langle W R^{\alpha 3} W \rangle$$

$$- \langle W R^{\alpha} W \rangle \langle W R^{\alpha 2} W \rangle.$$

21.

$$\begin{aligned} \textcircled{a} \quad |\tilde{\psi}^{(2)}\rangle &= (R^{(0)}W)^2 |\tilde{\psi}\rangle + \text{renorm. terms} \\ &= R^{(0)}W R^{(0)}W |\tilde{\psi}\rangle \\ &\quad - R^{(0)} \langle W \rangle R^{(0)}W |\tilde{\psi}\rangle \\ &= (R^{(0)}W)^2 |\tilde{\psi}\rangle - \langle W \rangle R^{(0)}W^2 |\tilde{\psi}\rangle, \end{aligned}$$

etc.

Back to MBPT.

3. Unperturbed and perturbed operators  
in MBPT.

We are interested in using RSPT to  
solve

$$K|\tilde{\psi}_0\rangle = k_0 |\tilde{\psi}_0\rangle, \quad (62)$$

where

$$K = H_N = H - \langle \tilde{\psi}_0 | H | \tilde{\psi}_0 \rangle \quad (63)$$

and

$$k_0 = \Delta E_0 = E_0 - \langle \tilde{\psi}_0 | H | \tilde{\psi}_0 \rangle.$$

$|F\rangle$  is the normalized FPM state defining the Fermi vacuum. Let us reorder the single-particle states such that the first  $N$  of them correspond to states occupied in  $|F\rangle$  (hole states) and single-particle states  $N+1, N+2, \dots$  are unoccupied (particle states). We will also use the standard notation for single-particle states:

- $i, j, \dots$  - hole states (occupied in  $|F\rangle$ )
- $a, b, \dots$  - particle states (unoccupied in  $|F\rangle$ )
- $p, q, \dots$  - generic states (occupied or unoccupied)

Thus,

$$i = 1, 2, 3, \dots, N$$

where  $N$  is the number of fermions in the system,  
and

$$a = N+1, N+2, \dots$$

With this notation,

$$|F\rangle = X_1^+ \dots X_N^+ |0\rangle = \prod_{i=1}^N X_i^+ |0\rangle$$

where  $|0\rangle$  is the true vacuum state.

### 3.1. Unperturbed problem.

We must define  $K_0$  and a single-particle basis such that

$$|\Psi\rangle = \prod_{i=1}^N X_i^\dagger |0\rangle \quad (64)$$

is an eigenstate of  $K_0$ . To do this, we recall that

[we could, in principle, use  $Z$  as an unperturbed operator, but then  $V$  is usually too big to obtain convergence of RSPT]

$$H = Z + V. \quad (65)$$

Let us approximate the two-body part of  $H$  by a one-body operator  $U$  and define

$$H_0 = Z + U. \quad (66)$$

where

$$U = \sum_{p,q} \langle p | \hat{u}(q) | q \rangle X_p^\dagger X_q \quad (\text{in 1st quantization}), \quad U = \sum_{i=1} \hat{u}(x_i),$$

↑ coordinates of fermion  $i$

We obtain

$$H_0 = \sum_{p,q} \langle p | Z + \hat{u}(q) | q \rangle X_p^\dagger X_q. \quad (67)$$

Let us further assume that  $\hat{u}$  is chosen such that we know how to solve the one-particle eigenvalue (or pseudoeigenvalue if  $\hat{u} = \hat{g} - \hat{z}$ , where  $\hat{f}$  is a Fock operator) problem,

$$(\hat{z} + \hat{u}) |p\rangle = \varepsilon_p |p\rangle. \quad (68)$$

With this choice of single-particle basis, we can write

$$H_0 = \sum_p \varepsilon_p X_p^+ X_p. \quad (69)$$

It is easy to show that any Slater determinant

$$|\Psi_{q_1 \dots q_N}\rangle = |\{q_1 \dots q_N\}\rangle \\ = X_{q_1}^+ X_{q_2}^+ \dots X_{q_N}^+ |0\rangle \quad (70)$$

is an eigenstate of  $H_0$  with an eigenvalue  $\varepsilon_{q_1} + \dots + \varepsilon_{q_N}$ . For example, in 1st quantization,

$$H_0 |\Psi_{q_1 \dots q_N}\rangle = \sum_{i=1}^N [\hat{z}(x_i) + \hat{u}(x_i)] A \left( \prod_{j=1}^N \chi_{q_j}(x_j) \right) \\ \stackrel{[A, H_0] = 0}{=} A \sum_{i=1}^N [\underbrace{\hat{z}(x_i) + \hat{u}(x_i)}_{\varepsilon_{q_i} \Psi_{q_i}(x_i)}] \chi_{q_i}(x_i) \\ \times \chi_{q_1}(x_1) \dots \chi_{q_{i-1}}(x_{i-1}) \chi_{q_{i+1}}(x_{i+1}) \dots \chi_{q_N}(x_N)$$

antisymmetrizer  $\frac{1}{N!} \sum_{\text{permutations}}$

$\langle r_1 | q_{r_1} \rangle$

$$\begin{aligned}
 &= A \left( \sum_{j=1}^N \varepsilon_{q_j} \right) \left( \prod_{n=1}^N \chi_{q_n}(x_n) \right) \\
 &= \left( \sum_{i=1}^N \varepsilon_{q_i} \right) |\Psi_{q_1 \dots q_N}\rangle. \quad (71)
 \end{aligned}$$

In particular,

$$H_0 |\Psi_0\rangle = E_0^{(0)} |\Psi_0\rangle,$$

where

$$E_0^{(0)} = \sum_{i=1}^N \varepsilon_i. \quad (72)$$

Let us then look at the remaining Slater determinants organized as particle-hole excitations from  $|\Psi_0\rangle$ . For example,

$$\begin{aligned}
 H_0 |\Psi_i^a\rangle &= (\varepsilon_1 + \dots + \varepsilon_{i-1} + \varepsilon_a + \varepsilon_{i+1} + \dots + \varepsilon_N) |\Psi_i^a\rangle \\
 &= [(\varepsilon_a - \varepsilon_i) + (\varepsilon_1 + \dots + \varepsilon_{i-1} + \varepsilon_i + \varepsilon_{i+1} + \dots + \varepsilon_N)] |\Psi_i^a\rangle \\
 &= [(\varepsilon_a - \varepsilon_i) + E_0^{(0)}] |\Psi_i^a\rangle. \quad (73)
 \end{aligned}$$

Similarly,

$$H_0 |\Psi_{ij}^{ab}\rangle = [(\varepsilon_a - \varepsilon_i + \varepsilon_b - \varepsilon_j) + E_0^{(0)}] |\Psi_{ij}^{ab}\rangle, \quad (74)$$

-26-

$$H_0 |\Psi_{i_1, \dots, i_n}^{\text{a, even}}\rangle = \left[ \sum_{j=1}^n (\varepsilon_{aj} - \varepsilon_{ij}) + E_0^{(0)} \right] |\Psi_{i_1, \dots, i_n}^{\text{a, even}}\rangle \quad (75)$$

Keeping the above in mind we define the unperturbed operator  $K_0$  used in MBPT as

$$\begin{aligned} K_0 &= H_0 - \langle \Psi_0 | H_0 | \Psi_0 \rangle \\ &= H_0 - E_0^{(0)}, \end{aligned} \quad (76)$$

where  $E^{(0)}$  is given by Eq. (72).

We obtain,

$$K_0 |\Psi_0\rangle = 0 = \chi_0(\Psi_0),$$

$$K_0 |\Psi_i^a\rangle = \chi_i^a |\Psi_i^a\rangle, \quad (77)$$

$$K_0 |\Psi_{i_1, \dots, i_n}^{\text{a, even}}\rangle = \chi_{i_1, \dots, i_n}^{\text{a, even}} |\Psi_{i_1, \dots, i_n}^{\text{a, even}}\rangle, \quad (77)$$

where

$$\chi_0 = 0, \quad (78)$$

$$\chi_i^a = \varepsilon_a - \varepsilon_i, \quad \dots$$

$$\chi_{i_1, \dots, i_n}^{\text{a, even}} = \sum_{j=1}^n (\varepsilon_{aj} - \varepsilon_{ij}), \quad n=1, \dots, N.$$

Determinants  $\langle \hat{E}_0 \rangle$ ,  $\langle \hat{E}_n^a \rangle$ , ... form over unperformed states  $\langle \hat{E}_n^i \rangle$  and  $\langle \hat{E}_n^b \rangle$ , ... form the corresponding unperformed eigenvalues  $\lambda_n^a, \lambda_n^b, \dots$

The many-body structure of  $K_0$  is

$$K_0 = Z + U - \langle \hat{E}_0 | Z + U | \hat{E}_0 \rangle \\ = Z_N + U_N, \text{ where} \quad (79)$$

$$Z_N = \sum_{pq} \langle \hat{e}_p | \hat{z} | \hat{e}_q \rangle N [X_p^+ X_q^-] \quad (80)$$

and

$$U_N = \sum_{pq} \langle \hat{e}_p | \hat{z} | \hat{e}_q \rangle N [X_p^+ X_q^-] \quad (81)$$

are the normal-ordered forms of  $Z$  and  $U$ , respectively. We can also write

$$K_0 = \sum_p \epsilon_p N [X_p^+ X_p^-]. \quad (82)$$

We recall that  $N[\dots]$  means move the (p-h) particle-hole creation operators ( $X_a^+, X_b^-$ ) to the left with respect to the corresponding p-h annihilation operators ( $X_a^-, X_b^+$ ) and multiply by the sign of the corresponding permutation needed for operator rearrangement.

-28-

$$Z = Z_N + \langle \hat{c}^\dagger \hat{c} \rangle, \text{ since, using}$$

Wick's theorem,

$$\begin{aligned} Z &= \sum_{pq} \langle p | \hat{c}^\dagger | q \rangle X_p^\dagger X_q \\ &= \sum_{pq} \langle p | \hat{c}^\dagger | q \rangle N[X_p^\dagger X_q] \\ &\quad + \sum_{pq} \langle p | \hat{c}^\dagger | q \rangle N[X_p^\dagger X_q] \\ &= Z_N + \sum_{pq} \langle p | \hat{c}^\dagger | q \rangle \chi(p) \delta_{pq}, \end{aligned} \tag{83}$$

where

$$\chi(p) = 1 \text{ if } p = i \text{ and } 0 \text{ if } p = \alpha.$$

(occupied p)    (unoccupied p)

This gives,

$$Z = Z_N + \sum_i \langle \hat{c}^\dagger \hat{c} | i \rangle = Z_N + \langle \hat{c}^\dagger \hat{c} \rangle \tag{84}$$

Similarly for  $U_N$  (and any one-body operator).

-29-

### 3.2. Perturbation

We want to write  $K = H_N$  as

$$K = K_0 + W.$$

Then,

$$\begin{aligned} W &= K - K_0 = (H - \langle \psi_0 | H | \psi_0 \rangle) \\ &\quad - (H_0 - \langle \psi_0 | H_0 | \psi_0 \rangle) \\ &= (H - H_0) - \langle \psi_0 | (H - H_0) | \psi_0 \rangle \\ &= \cancel{Z} + V - (\cancel{Z} + U) - \langle \psi_0 | \cancel{Z} + V - (\cancel{Z} + U) | \psi_0 \rangle \\ &= V - U - \langle \psi_0 | V - U | \psi_0 \rangle \quad (85) \\ &= V - \langle \psi_0 | V | \psi_0 \rangle - (U - \langle \psi_0 | U | \psi_0 \rangle) \end{aligned}$$

We already know that  $U - \langle \psi_0 | U | \psi_0 \rangle = U_N$ .

Using Wick's theorem, we can easily show that

$$V = \frac{1}{2} \sum_{pq,rs} \langle pq | \hat{v} | rs \rangle X_p^\dagger X_q^\dagger X_s X_r \quad (86)$$

$$= V_N + G_N + \langle \psi_0 | V | \psi_0 \rangle, \text{ where}$$

-30-

$$\begin{aligned} V_N &= \frac{1}{2} \sum_{pq,rs} \langle pq|\hat{\omega}|rs\rangle N [x_p^T x_q + x_q^T x_r] \\ &= \frac{1}{2} \sum_{pq,rs} \langle pq|\hat{\omega}|rs\rangle N [x_p^T x_q + x_q^T x_s] \\ &\quad (\text{or } \overset{1}{\cancel{q}} \dots \overset{1}{\cancel{q}} \langle pq|\hat{\omega}|rs\rangle_d \dots ), \end{aligned} \quad (87)$$

$$G_N = \sum_{pq} \langle p|\hat{g}|q\rangle N [x_p^T x_q], \quad (88)$$

with  $\langle p|\hat{g}|q\rangle = \sum_{i=1}^N \langle pi|\hat{\omega}|qi\rangle$

(mean field  
one-body potential  
creation ( $a_i^\dagger$ ) / destruction ( $a_i$ ))

$$\langle \hat{q}_0 | V | \hat{q}_0 \rangle = \frac{1}{2} \sum_{ij} \langle \hat{c}_j | \hat{s} | \hat{c}_i \rangle_A. \quad (89)$$

Thus,

$$W = V_N + G_N - U_N = W_1 + W_2, \quad (90)$$

where

and

$$W_1 = G_N - U_N = Q_N, \quad (91)$$

$$W_2 = V_N. \quad (92)$$

-3)-

Note that the one-body perturbation,

$$\begin{aligned} W_1 = Q_N &= \sum_{pq} \langle p | \hat{g} - \hat{u} | q \rangle N [x_p^\dagger x_q] \quad (93) \\ &= \sum_{pq} \langle p | (\hat{z} + \hat{g}) - (\hat{z} + \hat{u}) | q \rangle N [x_p^\dagger x_q] \\ &\quad \text{Fock operator } f \\ &= \sum_{pq} [K_p | \hat{f} | q \rangle - \epsilon_p \delta_{pq}] N [x_p^\dagger x_q], \end{aligned}$$

measures the departure of the single-particle basis from the Hartree-Fock case. Indeed, when  $|p\rangle$ 's are HF states,

$$W_1 = Q_N = 0, \text{ since in the HF case we use } \hat{u} = \hat{g} \text{ (or } (\hat{z} + \hat{u}) = \hat{f} \text{).}$$

In general though,

-32-

$$W_1 = Q_N = \sum_{r,s} \langle r|q|s\rangle N[X_r^{\dagger} X_s], \quad (94)$$

where  $\hat{q} = \hat{q} - \hat{u}$ , and

$$W_2 = \sqrt{N}. \quad (95)$$

### 3.3. Reduced resolvent in MBPT

We know that  $R^{(0)} = \sum_{n>0} \frac{|\bar{q}_n\rangle\langle\bar{q}_n|}{\epsilon_0 - \epsilon_n}$ .

In our case,  $|\bar{q}_n\rangle$ 's are  $|\bar{q}_0\rangle, |\bar{q}_1\rangle, \dots$

Thus,

$$R^{(0)} = \sum_{n=1}^N \sum_{\substack{i_1 < \dots < i_n \\ q_1 < \dots < q_n}} \frac{|\bar{q}_{i_1, \dots, i_n}^{(0)}\rangle\langle\bar{q}_{i_1, \dots, i_n}^{(0)}|}{\epsilon_0 - \epsilon_{i_1, \dots, i_n}^{(0)}}, \quad (96)$$

$$\text{where } \epsilon_0 = 0 \text{ and } \epsilon_{i_1, \dots, i_n}^{(0)} = \sum_{j=1}^n (\epsilon_{q_j} - \epsilon_{i_j}).$$

This allows us to write

$$R^{(0)} = \sum_{n=1}^N R_n^{(0)}, \quad (97)$$

where the n-body component of  $R^{(0)}$  is

-33-

$$R_n^{(0)} = \sum_{\substack{i_1 < \dots < i_n \\ a_1 < \dots < a_n}} \frac{|\langle \tilde{\psi}_{i_1 \dots i_n}^{a_1 \dots a_n} \rangle \times \langle \tilde{\psi}_{i_1 \dots i_n}^{a_1 \dots a_n} |}{\omega_{i_1 \dots i_n}^{a_1 \dots a_n}}$$
$$= \left(\frac{1}{n!}\right)^2 \sum_{\substack{i_1 \neq \dots \neq i_n \\ a_1 \neq a_2 \dots \neq a_n}} \frac{|\langle \tilde{\psi}_{i_1 \dots i_n}^{a_1 \dots a_n} \rangle \times \langle \tilde{\psi}_{i_1 \dots i_n}^{a_1 \dots a_n} |}{\omega_{i_1 \dots i_n}^{a_1 \dots a_n}}$$

ALLOWING  
EPV determinants  
which are zero here

$$\left(\frac{1}{n!}\right)^2 \sum_{\substack{i_1 \dots i_n \\ a_1 \dots a_n}} \frac{|\langle \tilde{\psi}_{i_1 \dots i_n}^{a_1 \dots a_n} \rangle \times \langle \tilde{\psi}_{i_1 \dots i_n}^{a_1 \dots a_n} |}{\omega_{i_1 \dots i_n}^{a_1 \dots a_n}}$$

MBPT Denominator (98)

with

$$\omega_{i_1 \dots i_n}^{a_1 \dots a_n} = \sum_{g=1}^n (\epsilon_{ig} - \epsilon_{og}), \quad (99)$$

$$\langle \tilde{\psi}_{i_1 \dots i_n}^{a_1 \dots a_n} | = \prod_{g=1}^n X_{ag}^+ X_{ig}, \quad (100)$$

$$\langle \tilde{\psi}_{i_1 \dots i_n}^{a_1 \dots a_n} | = (\langle \tilde{\psi}_{i_1 \dots i_n}^{a_1 \dots a_n} |)^+ = \prod_{g=1}^n X_{ig}^+ X_{ag}$$

In RSP, we need powers of  $R^{(0)}$  as well,

-34-

$$(R^{(0)})^k = \sum_{n>0} \frac{[\bar{\epsilon}_n \sum d_n]}{(\bar{\epsilon}_0 - \bar{\epsilon}_n)^k}, \quad (102)$$

because of orthonormality of  $P_{\bar{\epsilon}_n}$ 's,  
all that changes is power of  $\bar{\epsilon}_0 - \bar{\epsilon}_n$

In our case, because of orthonormality of Slater determinants,

$$(R^{(0)})^k = \sum_{n=1}^N (R_n^{(0)})^k, \quad (103)$$

where

$$(R_n^{(0)})^k = \left(\frac{1}{n!}\right)^2 \sum_{\text{spins}} \frac{E_{\text{cyclic}}^{\text{even}} [\bar{\epsilon}_0 \times \bar{\epsilon}_1 \times \dots \times \bar{\epsilon}_{n-1}]}{(c\omega_{\text{cyclic}})^k} \quad (104)$$

$\nearrow$   
k-th power  
of the MBPT  
denominator

### 3.4. MBPT energy and wave function corrections.

We know that  $K = K_0 + W$  ( $K = E_N$ ),

$$K_0 = Z_N + U_N \text{ and } W = W_1 + W_2,$$

where  $W_1 = Q_N$  and  $W_2 = U_N$  are both in

the normal product form. Because of the latter  
observation,

$$\langle \bar{\phi}_0 | N [ \dots ] \bar{\phi}_0 \rangle = 0$$

$$k_0^{(1)} = \langle \bar{\phi}_0 | W(\bar{\phi}_0) \rangle = 0. \quad (105).$$

This simplifies the MBPT analysis using

$K = H_N$ . We obtain ( $\langle \dots \rangle$  means  $\langle \bar{\phi}_0 | \dots | \bar{\phi}_0 \rangle$ ),

$$\Delta E_f^{(1)} \equiv k_0^{(1)} = \langle \bar{\phi}_0 | W(\bar{\phi}_0) \rangle = \langle W \rangle = 0,$$

$$\Delta E_f^{(2)} \equiv k_0^{(2)} = \langle W R^{(0)} W \rangle,$$

$$\Delta E_f^{(3)} \equiv k_0^{(3)} = \langle W (R^{(0)} W)^2 \rangle - \langle W \rangle \langle W R^{(0)} W \rangle \\ \stackrel{\text{def}}{=} \langle W (R^{(0)} W)^2 \rangle \quad \leftarrow \text{no renormalization terms yet!}$$

The 1st  
occurrence  
of renormaliza-  
tion terms

$$\rightarrow k_0^{(4)} \stackrel{\text{after eliminating the } k_0^{(3)} \text{ terms}}{=} \langle W (R^{(0)} W)^3 \rangle - \langle W R^{(0)} W \rangle \\ \times \langle W R^{(0)} W \rangle,$$

etc.

(106)

IMPORTANT FOR

DIAGRAM CANCELLATION ANALYSIS THAT LEADS TO LINKED CLUSTER THEORY

-36-

Similarly,

$$|\bar{\psi}^{(1)}\rangle = R^{(0)}W|\bar{\phi}\rangle,$$

$$\begin{aligned} |\bar{\psi}^{(2)}\rangle &= (R^{(0)}W)^2|\bar{\phi}\rangle - \langle W \rangle R^{(0)2}W|\bar{\phi}\rangle \\ &= (R^{(0)}W)^2|\bar{\phi}\rangle, \end{aligned}$$

$$|\bar{\psi}^{(3)}\rangle = (R^{(0)}W)^3|\bar{\phi}\rangle - \langle WR^{(0)}W \rangle R^{(0)2}W|\bar{\phi}\rangle,$$

etc.

*1st occurrence of renormalization terms*

(important for linked cluster theorem)

(107)

Finally,

$$k_0 = \Delta E_0 = E_0 - \langle \bar{\phi}_0 | H(\bar{\phi}_0) \rangle \quad (108)$$

$$= \chi_0 + k_0^{(1)} + k_0^{(2)} + \dots$$

$$= k_0^{(2)} + \dots = \sum_{n=2}^{\infty} k_0^{(n)} = \sum_{n=2}^{\infty} \Delta E_0^{(n)}$$

$\chi_0 = 0$   
 $k_0^{(1)} = 0$

Correlation energy starts in the second order.

This is easy to understand. If we used  $H$  rather than  $H^N$  and  $H = Z + U$  neither, then the shifted  $K_0 = H_0 - \langle \psi_0 | H_0 | \psi_0 \rangle$ , we would have

$$K = H = \tilde{K}_0 + \tilde{W}, \text{ where}$$

$$\tilde{K}_0 = Z + U \text{ and } \tilde{W} = V - U.$$

In that case, the energy  $E_0$  ( $\in H(\psi_0) = E_0 | \psi_0 \rangle$ ) would become

$$E_0 = E_0^{(0)} + E_0^{(1)} + E_0^{(2)} + \dots,$$

where

$$E_0^{(0)} = \langle \psi_0 | \underbrace{Z + U}_{\tilde{K}_0} | \psi_0 \rangle = \sum_{i=1}^N \varepsilon_i,$$

$$E_0^{(1)} = \langle \psi_0 | V - U | \psi_0 \rangle = \langle \psi_0 | \tilde{W} | \psi_0 \rangle$$

etc. Now,

$$E_0^{(0)} + E_0^{(1)} = \langle \psi_0 | \tilde{K}_0 + \tilde{W} | \psi_0 \rangle$$

$$= \langle \psi_0 | (Z + U) + (V - U) | \psi_0 \rangle$$

$$= \langle \psi_0 | H | \psi_0 \rangle \Rightarrow \text{mean-field energy, no correlations!}$$

-<sup>38</sup> reference energy correction.

Thus,

$$E_0 = \underbrace{\langle \tilde{\psi}_0 | H(\tilde{\psi}_0) \rangle}_{\text{0th + 1st order}} + \underbrace{\tilde{E}_0^{(2)} + \dots}_{\text{2nd order}}$$

#### 4. Diagrammatic representation of MBPT energy and wave function corrections.

In order to evaluate MBPT expressions for the energy and wave function corrections, we need to evaluate quantities of the following types:

$$\langle \tilde{\psi}_0 | W(R^{(0)})^{n_1} W(R^{(0)})^{n_2} W_{\dots} (R^{(0)})^{n_y} \delta W | \tilde{\psi}_0 \rangle \quad (109)$$

(energy corrections)

$$(R^{(0)})^{n_1} W(R^{(0)})^{n_2} W_{\dots} (R^{(0)})^{n_y} \delta W | \tilde{\psi}_0 \rangle \quad (110)$$

(wave function corrections),

where  $n_1, n_2, \dots, n_y \geq 1$  and  $W = W_1 + W_2$ .

If we want to do it diagrammatically, we must come up with a diagrammatic representation of  $W_1$ ,  $W_2$ , and  $R^{(0)K}$  and then follow the diagrammatic rules to determine the final formulas.

## QUICK SUMMARY OF DIAGRAMMATIC CALCULATIONS

Diagrammatic formalism allows us to calculate expressions of the following form:

$$K_A \dots K_Z = \sum_C K_C, \quad (\text{III})$$

where each  $K_C$  is a many-body operator in the standard or normal product form. We could, of course, do this algebraically, using Wick's theorem, but Wick's theorem often produces multiple copies of the same term, which we have to manually recognize and collect. Diagrams are nicer in this regard, since one only has to determine the non-equivalent admissible resulting diagrams, relevant to the problem of interest. All of the redundancies are taken care of by the so-called topological weights. Here, because of time constraints, we will focus on Husenholtz diagrams, which are used to represent many-body operators employing

-40-

antisymmetrized matrix elements. For example  
the  $\hat{O}_k$ -body operator in the normal-product  
form

$$\begin{aligned}\hat{O}_k &= \left(\frac{1}{k!}\right)^2 \sum_{\substack{p_1, \dots, p_k \\ q_1, \dots, q_k}} \langle p_1 \dots p_k | \hat{O}_k | q_1 \dots q_k \rangle_A \\ &\quad \times N [X_{p_1}^+ \dots X_{p_k}^+ X_{q_1}^- \dots X_{q_k}^-] \\ &= \left(\frac{1}{k!}\right)^2 \sum_{\substack{p_1, \dots, p_k \\ q_1, \dots, q_k}} \langle p_1 \dots p_k | \hat{O}_k | q_1 \dots q_k \rangle_A \\ &\quad \times N [X_{p_1}^+ X_{q_1}^- \dots X_{p_k}^+ X_{q_k}^-], \quad (112)\end{aligned}$$

where

$$\begin{aligned}\langle p_1 \dots p_k | \hat{O}_k | q_1 \dots q_k \rangle_A &= \sum_{R \in S_k} (-1)^R \\ &\quad \times \langle p_1 \dots p_k | \hat{O}_k | q_{R_1} \dots q_{R_k} \rangle_A, \quad (113)\end{aligned}$$


with

$$R = \begin{pmatrix} 1 & \dots & k \\ R_1 & \dots & R_k \end{pmatrix} \quad (114)$$

representing the index permutation, is represented by

-41-

a vertex



Outgoing lines are  $X^+$ , incoming lines are  $X$ ,  
 $(\frac{1}{k!})^2$

factor is taken care of by the equivalences  
among fermion lines  $p_1, \dots, p_k$  and  $q_1, \dots, q_k$ .  
The vertex is always drawn in a way  
specific to the operator of interest. We will show  
the  $W_1, W_2$ , and  $(R_n^{(0)})^\dagger$  operators diagrammatically  
in a moment.

Once we represent operators  $K_A, \dots, K_Z$  on the  
left-hand side of Eq. (III), we proceed as  
follows (we will assume that all fermion lines carry free  
labels which are summed over):


- (1) Draw the nonoriented (no arrows) Hugenholtz  
skeletons corresponding to  $K_A, \dots, K_Z$  along the  
fictitious time line ( $\alpha$  is), going in this prescript by  
from left to right, and form the non-equivalent  
resulting Hugenholtz skeletons (or their  
subset relevant to the calculation of interest) of below

Formation of the resulting diagrams is accomplished by connecting fermion lines. Such connections represent contractions of  $X$  and  $X^\dagger$  operators, as in Wick's theorem,

$$\begin{aligned}
 M_1 \dots M_m &= N[M_1 \dots M_m] + \sum_{\mu < \nu} N[M_1 \dots \overset{\mu}{M} \dots M_\nu \dots M_m] \\
 &+ \sum_{\substack{\mu_1 < \nu_1, \\ \mu_2 < \nu_2, \\ \mu_1 < \mu_2, \\ \nu_1 \neq \nu_2}} N[M_1 \dots \overset{\mu_1}{M} \dots M_{\mu_2} \dots \overset{\nu_1}{M} \dots M_{\nu_2} \dots M_m] \\
 &+ \dots \quad (116)
 \end{aligned}$$

- (2) Add arrows to fermion lines in all possible allowed ways (for example,  $k$  lines have arrows toward  $\otimes$  and  $k$  lines leave  $\otimes$  in (115)). Lines that remain unconnected must carry the same orientation as on the left-hand side of Eq. (111), unless a particular expression forces a modification.

- (3) Add the appropriate spin-orbital or single-particle indices to each line in the resulting Hugenholtz diagrams. For the internal lines going from left to right, use hole indices, e.g.,



$$(X_i^\dagger X_j = \delta_{ij})$$

For the internal lines going from right to left, use particle indices, e.g.,

$$\xleftarrow{a} \quad (X_a^a X_b^f = \delta_{ab})$$

Uncontracted external fermion lines retain their character from the left-hand side of Eq. (II), unless the actual expression forces some adjustment.

For example, in MBPT, all external lines will extend to the left, since  $X_a |\Psi_0\rangle = X_a^f |\Psi_0\rangle = 0$ , and the normal ordering places  $X_a^a$  and  $X_i^f$  in the rightmost positions, allowing direct action on  $|\Psi_0\rangle$ . In MBPT, operator products always act on  $|\Psi_0\rangle$ . In other words, in MBPT we can only have external lines of the following two types:

$$\begin{array}{c} \xleftarrow{a} \\ \xleftarrow{i} \end{array} = X_a^f$$

As shown below,  $R^{(0)}$  in wave function corrections enforces the same.

(4) Read the resulting Hugenholtz diagrams.


The last step is executed as follows:

(a) Determine the topological weight

$$W_R^{(H)}$$

based on the equivalences among fermion lines  
in the oriented Hugenholtz & de Jong  
(diagram stripped of free choices labeling  
fermion lines) corresponding to a given  
resulting diagram  $R$ .

(b) Draw the Brandon representative (one of  
the Goldstone diagrams corresponding to  
Hugenholtz diagram  $R$ ) by expanding  
Hugenholtz vertices to their Goldstone-like  
form, e.g.



Assign the scalar factor  $d_{X_R^1 X_R^2}^{(B)}$   
being a product of  
antisymmetrized matrix elements,  
labeled by appropriate internal ( $X_R^1$ )  
and external ( $X_R^2$ ) indices at various  
fermion lines, as seen in the Brandon  
diagram representing  $R$ .

-45-

$l_R^{(B)} + h_R^{(B)}$   $\downarrow$   $s_R^{(B)}$  (sign factor)

Assign the sign  $(-1)^{l_R^{(B)} + h_R^{(B)}}$  where  $l_R^{(B)}$  and  $h_R^{(B)}$  are the numbers of closest loops and internal hole lines in the Bremsstrahlung diagram.

Assign, if relevant, the operator expression

$$\hat{O}_R^{(B)} = N \left[ \prod_{r=1}^{m_R^{(B)}} X_p^r X_q^r \right]$$

to the diagram, where  $X_p^r$  and  $X_q^r$  correspond to external lines  $p_r$  and  $q_r$  exiting and entering open path  $r$  in Bremsstrahlung diagram  $R$ . ( $m_R^{(B)}$  is the total number of open paths). In MBPT,  $p_r$  must be a particle line and  $q_r$  must be a hole line, as explained above. The final formula for  $K_A \dots K_Z$  is

$$K_A \dots K_Z = \sum_R K_R , \quad (17)$$

where the summation on the right-hand side involves only the non-equivalent resulting diagrams (relevant to the problem of interest; cf. below) and

$$K_R = (-1)^{l_R^{(B)} + h_R^{(B)}} W_R^{(H)} \sum_{X_p^r X_q^r} \hat{O}_R^{(B)}$$

indices of internal lines

In MBPT, we only have two situations:

- energy diagrams that correspond to Eq. (109), meaning resulting diagrams with no external lines, so that

$$K_R = (-1)^{l_R^{(B)} + h_R^{(B)}} w_R^{(H)} \sum_{X'_R} d_{X'_R}^{(B)}. \quad (118)$$

- wave function diagrams that correspond to Eq. (110), where all external lines extend to the left, as in  $\overleftarrow{\bullet} = X_a^+$  and  $\overrightarrow{\bullet} = X_c^-$ , so that


$$K_R = (-1)^{l_R^{(B)} + h_R^{(B)}} w_R^{(H)}$$

indices of  
external lines in  
open paths

$$\times \sum_{X'_R(a_1, \dots, a_m, c_1, \dots, c_m)} d_{X'_R(a_1, \dots, a_m, c_1, \dots, c_m)}^{(B)} \\ \times N [X_{a_1}^+ X_{c_1}^- \dots X_{a_m}^+ X_{c_m}^-] (\bar{e})$$

indices of internal lines

(119)



Branckow diagram R.  
(120)


The requirement that all external lines must extend to the left is also enforced by the presence of the freezed resolvent in the leftmost position of Eq. (110).

Now, we introduce Hugenholtz and Branckow vertices representing  $W_1 = Q_N$ ,  $W_2 = V_N$ , and  $R_n^{(0)}$ :

- $W_1 = Q_N = \sum_{r,s} \langle r | \hat{q} | s \rangle N [X_r^+ X_s^-]$ , where

$$\hat{q} = q - u.$$

Hugenholtz  
and Branckow  
look identical  
since  $W_1$  is  
one-body



$$W_{Q_N}^{(H)} = 1$$

outgoing

$$d_{rs}^{(B)} = \langle r | \hat{q} | s \rangle$$

incoming

$$\hat{O}_{rs}^{(B)} = N [X_r^+ X_s^-]$$

-48-


$$W_1 = w_{V_N}^{(H)} \sum_{rs} d_{rs}^{(B)} \hat{O}_{rs}^{(B)}$$

$(l_{V_N}^{(B)} = 0)$   
 $h_{V_N}^{(B)} = 0$

$$\bullet \quad W_2 = V_N = \frac{1}{4} \sum_{rstu} \langle r s | \hat{o} | t u \rangle$$


$\frac{1}{2})^2$

$$\times N [X_r^+ X_t^+ X_s^+ X_u^-]$$



Hugenholtz

$\Rightarrow$



Brandow (Goldstone representative)

$$W_{V_N}^{(H)} = \frac{1}{4} \downarrow \downarrow \text{outgoing} \quad \downarrow \downarrow \text{incoming}$$

$$d_{rstu}^{(B)} = \langle r s | \hat{o} | t u \rangle$$

$$\hat{O}_{rstu}^{(B)} = N [ \underbrace{X_r^+ X_t^+}_{\text{open path}} \underbrace{X_s^+ X_u^-}_{\text{open path}} ]$$

$$W_2 = w_{V_N}^{(H)} \sum_{rstu} d_{rstu}^{(B)} \hat{O}_{rstu}^{(B)}$$

$(l_{V_N}^{(B)} = 0)$   
 $h_{V_N}^{(B)} = 0$

- Reduced resolvent, focus on k-th power of the n-body component  $(R_n^{(0)})^k$ .


$$(R_n^{(0)})^k = \left(\frac{1}{n!}\right)^2 \sum_{\substack{\text{Circin} \\ \text{Circin}}} \frac{E_{\text{Circin}}^{\text{Circin}} |\langle \Phi \rangle \times \langle \Phi \rangle| E_{\text{Circin}}^{\text{Circin}}}{(c_{\text{Circin}}^{\text{Circin}})^k}$$

$$E_{\text{Circin}}^{\text{Circin}} = N [X_{a_1}^+ X_{i_1}^- \dots X_{a_n}^+ X_{i_n}^-]$$

$$E_{\text{Circin}}^{\text{Circin}} = (E_{\text{Circin}}^{\text{Circin}})^+ = N [X_{a_1}^+ X_{i_1}^- \dots X_{i_n}^+ X_{a_n}^-]$$

$$\omega_{\text{Circin}}^{\text{Circin}} = -\delta_{\text{Circin}}^{\text{Circin}} = \sum_{g=1}^N (\epsilon_i^g - \epsilon_a^g)$$

↑  
Circin hole particle  
↓  
 $\sum_{a=1}^N$



$$W_{R_n^{(0)}}^{(H)} = \left(\frac{1}{h_0}\right)^2$$

$$d_{c_{1, \dots, i, \dots, n}, d_{1, \dots, n}}^{(B)} = (\omega_{c_{1, \dots, i, \dots, n}}^{d_{1, \dots, n}})^{-k}$$

$$= \left[ \sum_{g=1}^n (\varepsilon_{ig} - \varepsilon_{og}) \right]^{-k}$$

lines "sliced" by !

$$\hat{O}_{c_{1, \dots, i, \dots, n}, d_{1, \dots, n}}^{(B)} = E_{c_{1, \dots, i, \dots, n}}^{d_{1, \dots, n}} | \Xi_0 \times_{\delta_0} | E_{d_{1, \dots, n}}^{c_{1, \dots, i, \dots, n}}$$

$$(R_n^{(0)})^k = W_{R_n^{(0)}}^{(H)} \sum_{\substack{c_{1, \dots, i, \dots, n} \\ d_{1, \dots, n}}} d_{c_{1, \dots, i, \dots, n}, d_{1, \dots, n}}^{(B)} \hat{O}_{c_{1, \dots, i, \dots, n}, d_{1, \dots, n}}^{(B)}$$

We can see how the  $R_n^{(0)}$  in the leftmost position in wave function expression enforces the requirement that external lines extend to the left representing


$$E_{c_{1, \dots, i, \dots, n}}^{d_{1, \dots, n}} = N \prod_{g=1}^n \times_{\delta_g}^+ X_{ig} ] \text{ acting on } |\Xi_0 \rangle.$$

-51-

Equipped with the above information, let us examine the second-order correction to energy,


$$\begin{aligned}
 k_0^{(2)} &= \langle \psi_0 | W R^{(0)} W | \psi_0 \rangle \\
 &= \langle \psi_0 | Q_N R^{(0)} Q_N | \psi_0 \rangle + \langle \psi_0 | Q_N R^{(0)} | \psi_0 \rangle \\
 &\quad + \langle \psi_0 | V_N R^{(0)} Q_N | \psi_0 \rangle + \langle \psi_0 | V_N R^{(0)} V_N | \psi_0 \rangle \\
 &= k_0^{(2)}(A) + k_0^{(2)}(\cancel{X}_1) + k_0^{(2)}(\cancel{X}_2) + k_0^{(2)}(B)
 \end{aligned} \tag{21}$$

$k_0^{(2)}(A)$  in Hohenberg representation ( $R^{(0)} = \sum_{n=1}^N R_n^{(0)}$ ):



one  
cannot  
contract  
lines  
on  
 $Q_N$   
due to  
normal  
orderings  
or a lines  
within lines

Thus, we must ~~contract~~ (connect) p and q lines with  $a_1, \dots, a_n, i_1, \dots, i_n$  in lines to the left of the dashed 'slicing' line. Similarly, r and s must be connected to lines  $a_1, \dots, a_n, q_1, \dots, q_n$  extending to the right relative to the dashed slicing line. This can only be done when  $n = 1$ ! We obtain



1 (or 2)      1 (or 2)

$$\text{sign factor } // \quad //$$

$$S_A^{(B)} = (-1)^{L_A^{(B)} + h_A^{(B)}} = +1,$$


$$d_{ia}^{(B)} = \langle i | \hat{q} | a \rangle \langle a | \hat{q} | i \rangle (\varepsilon_i - \varepsilon_a)^{-1}$$

$(\omega_i^a)^{-1}$

Thus,

$$K_0^{(2)}(A) = \sum_{i,a} \frac{\langle i | \hat{q} | a \rangle \langle a | \hat{q} | i \rangle}{\varepsilon_i - \varepsilon_a} \quad (122)$$

Please note that we could obtain this result by drawing



or even




obtained from  $Q_N Q_N$  only if we agreed on

an additional denominator convention that with each pair of neighboring W ( $Q_N$  or  $V_N$ ) vertices we associate the energy denominator obtained by assigning

$$[(\varepsilon_{i_1} - \varepsilon_{a_1}) + \dots + (\varepsilon_{i_n} - \varepsilon_{a_n})]^{-k}$$

to lines



in the region


between the neighboring Ws where there is  $[R_n^{(0)}]^k$ .

Let us examine the remaining contributions to  $K^{(2)}$ .

$$K_0^{(2)}(X)$$



$$R_n^{(0)}$$




$$\langle \psi_1 | \dots | \psi_n \rangle \langle \psi_n | \dots | \psi_1 \rangle$$

Again, we must connect lines  $p, q$  with lines  $a_1, i_1, \dots, a_n, i_n$  extending to the left of the dashed line slicing the reduced resolvent. We also must fully contract lines  $r, s, t, u$  with lines  $a_1, i_1, \dots, a_n, i_n$  extending to the right of the slicing dashed line. The former means  $n=1$ . The latter  $n=2$ . We cannot have it both ways,

so

$$k_0^{(2)}(X) = 0. \quad (123)$$

Note that we do not need diagrams representing  $R^{(0)}$  to come up with such a result since we cannot produce a diagram without external lines from




$\mathcal{Q}_N$

$V_N$

(we are not allowed to contract lines on  $V_N$  since  $V_N$  is in the normal ordered form  $\rightarrow$   $\rightarrow$  generalized Wick's theorem).

Similarly,


$$k_0^{(2)}(X_2) = 0. \quad (124)$$



We must fully contract lines between  $\langle \Phi_0 |$  (or  $| \Phi_0 \rangle$ ) and  $| \Phi_0 \rangle$  and between  $\langle \Phi_0 |$  (or  $| \Phi_0 \rangle$ ) and  $\langle \Phi_0 |$ .

This is only possible when  $n=2$  (we cannot contract lines on the same  $V_N$  due to normal ordering).


We obtain :



①

(Hugenholtz diagram)

The corresponding Brandon diagram looks as follows:



We obtain,

$$W_B^{(H)} = \frac{1}{4} \quad (\text{a, b equivalent; i, j equivalent}),$$

$$S_B^{(B)} = (-)^{l_B^{(B)} + h_B^{(B)}} = +1, \quad (l_B^{(B)} = 2 \text{ or } 4, h_B^{(B)} = 2 \text{ or } 4)$$

$$d_{ijab}^{(B)} = \langle ij|\hat{o}|ab\rangle_A \langle ab|\hat{o}|ij\rangle_A$$

$$\times \left( \varepsilon_i - \varepsilon_a + \varepsilon_j - \varepsilon_b \right)^{-1},$$

$$(o_{ij})^{-1}$$


$$K_0^{(2)}(B) = \frac{1}{4} \sum_{ijab} \frac{\langle ij|\hat{o}|ab\rangle_A \langle ab|\hat{o}|ij\rangle_A}{\varepsilon_i - \varepsilon_a + \varepsilon_j - \varepsilon_b} \quad (125)$$

-57-

Once again, the only role of the reduced resolvent is to introduce the denominator

$$(\varepsilon_i - \varepsilon_a + \varepsilon_j - \varepsilon_b)^{-1},$$

obtained by slicing the lines between  $V_N$  vertices in a Huyghen's diagram obtained for  $V_N \cdot V_N$ ,




(126)

Line  $i$  going from left to right contributes  $\varepsilon_i$ , line  $a$  going from right to left contributes  $-\varepsilon_a$ , for a total of  $(\varepsilon_i - \varepsilon_a)$  contribution for this pair of lines. The rest of the expression, i.e.,

$$\frac{1}{4} \langle i j | \hat{\sigma} ab \rangle_A \langle ab | \hat{\sigma} ij \rangle_A$$

can be read from diagram (126) and its Brauer counterpart,




-88-

We see similar patterns in wave function expressions in first order,

$$\begin{aligned} |\psi_0^{(1)}\rangle &= R^{(0)} W |\psi_0\rangle \\ &= R^{(0)} Q_N |\psi_0\rangle + R^{(0)} V_N |\psi_0\rangle \\ &= |\psi_0^{(1)}(A)\rangle + |\psi_0^{(1)}(B)\rangle. \end{aligned} \quad (27)$$

$|\psi_0^{(1)}(A)\rangle$ :



all lines must be fully contracted between  $|\psi_0\rangle$  and  $|\psi_0^{(1)}\rangle$ .  
This means that  $n=1$ .

We obtain,


$$= \sum_{i,a} \frac{\langle \alpha_i | \bar{a}_i \rangle}{\epsilon_i - \epsilon_a} |\bar{a}_i\rangle$$

$\omega_i^a$

$N[\bar{a}_i | \bar{a}_i] \langle \bar{a}_i |$

(128)

We could obtain this from



if we adopted the additional convention for lines sliced by the resolvent line.

Similarly,

$$|\Psi_0^{(1)}(B)\rangle = R_n^{(0)} V_N |\bar{a}_i\rangle$$

$\sum_{n=1}^N \langle \bar{a}_i | \bar{a}_i \rangle$


$\langle \bar{a}_i | \bar{a}_i \rangle = \langle \bar{a}_i | \bar{a}_i \rangle \langle \bar{a}_i | \bar{a}_i \rangle$

to connect fully here,  $n$  must be 2

-60-

We obtain

$$E_i^a E_j^b | \tilde{\psi} \rangle = E_{ij}^{ab} | \tilde{\psi} \rangle = | \tilde{\psi}_{ij}^{ab} \rangle$$




Hugenholtz

Brandonon

$$= \frac{1}{4} \sum_{cijab} \frac{\langle ab | \tilde{\psi} | ij \rangle_A}{\varepsilon_i - \varepsilon_a + \varepsilon_j - \varepsilon_b} | \tilde{\psi}_{ij}^{ab} \rangle. \quad (129)$$

$\tilde{\psi}_{ij}^{ab}$

We could obtain this from a Hugenholtz diagram



If we adopted the denominator convention,

The denominator convention of rescaling denominators from lines sliced between the neighboring  $W$ s and the external lines originates from the observation that, in general, every MBPT expression has a structure


$$\dots W (R^{(0)})^m W (R^{(0)})^n W \dots$$

$Q_{N \text{ or } V_N}$

$Q_N \text{ or } V_N$

$Q_{N \text{ or } V_N}$

or, diagrammatically (schematically),



(between  $\langle \$ \rangle$  and  $\langle \$ \rangle$ )


or between  $(R^{(0)})^m$  and  $(R^{(0)})^n$ )

MUST BE FULLY CONTRACTED

gives  
MBPT  
denominator

In other words, we do not need to draw diagrams representing reduced resolvents and can use the standard rules of constructing the resetting energy and wave function corrections from  $\mathcal{W}$  vertices only, as if  $R^0$ 's were not present. If we adopt the denominator convention, i.e. the incorporation of denominators corresponding to fermion lines between the neighboring  $W$  vertices with the powers corresponding to powers of resolvents between these  $W$ s. The energy diagrams have no external lines and the wave function diagrams have lines extending to the left (representing excited determinants).

The leftmost external lines in the wave function diagrams are also accompanied by the denominators corresponding to  $(R^0)^k$  showing up in the leftmost position in  $\langle \mathcal{W} \rangle^0$ . The denominator convention automatically excludes diagrams with "dangerous denominators" where there are no lines between neighboring  $W$ s,



which would formally result from a singlet

-63-

expression  
that

$$\frac{|\psi_0\rangle\langle\psi_0|}{\epsilon_0 - \epsilon_0}$$


We have learned

$$\Delta E_0^{(2)} = K_0^{(2)} = k_0^{(2)}(A) + k_0^{(2)}(B), \quad (130)$$

where

$$k_0^{(2)}(A) = \langle \psi_0 | Q_N R^{(2)} Q_N | \psi_0 \rangle$$

$$= \langle \psi_0 | Q_N R^{(2)}_1 Q_N | \psi_0 \rangle$$





$$= \sum_{i,a} \frac{\langle i | \hat{q} | a \rangle \langle a | \hat{q} | i \rangle}{\epsilon_i - \epsilon_a}, \quad (131)$$

$$k_0^{(2)}(B) = \langle \psi_0 | V_N R^{(2)} V_N | \psi_0 \rangle$$

$$= \langle \psi_0 | V_N R^{(2)}_2 V_N | \psi_0 \rangle$$

=



Hugenholtz

Brendow

-64-

$$= \frac{1}{4} \sum_{cjab} \frac{\langle c_j | \hat{o} | ab \rangle_A \langle ab | \hat{o} | c_j \rangle_A}{\varepsilon_c - \varepsilon_a + \varepsilon_j - \varepsilon_b}$$

$$= \frac{1}{2} \sum_{cjab} \frac{\langle c_j | \hat{o} | ab \rangle \langle ab | \hat{o} | c_j \rangle_A}{\varepsilon_c - \varepsilon_a + \varepsilon_j - \varepsilon_b} . \quad (B2)$$

$$|\Psi_0^{(1)}\rangle = |\Psi_0^{(1)}(A)\rangle + |\Psi_0^{(1)}(B)\rangle, \quad (B3)$$

where

$$|\Psi_0^{(1)}(A)\rangle = R^{(0)} Q_N |\bar{\psi}_0\rangle = R_1^{(0)} Q_N |\bar{\psi}_0\rangle$$

$$= \begin{array}{c} a \\ \diagup \\ c \\ \diagdown \\ i \\ \diagup \\ b \\ \diagdown \\ j \\ \textcircled{1} \end{array} = \sum_{ca} \frac{\langle a | \hat{o} | i \rangle}{\varepsilon_i - \varepsilon_a} |\bar{\psi}_i^a\rangle, \quad (B4)$$

$$|\Psi_0^{(1)}(B)\rangle = R_2^{(0)} V_N |\bar{\psi}_0\rangle = R_2^{(0)} V_2 N |\bar{\psi}_0\rangle$$

$$= \begin{array}{c} a \\ \diagup \\ c \\ \diagdown \\ i \\ \diagup \\ b \\ \diagdown \\ j \\ \textcircled{1} \end{array} \quad \text{or} \quad \begin{array}{c} a \\ \diagup \\ i \\ \diagdown \\ b \\ \diagup \\ j \\ \textcircled{1} \end{array} \quad (B5)$$

Hugenholtz Brandon

$$= \frac{1}{4} \sum_{cjab} \frac{\langle ab | \hat{o} | c_j \rangle_A}{\varepsilon_c - \varepsilon_a + \varepsilon_j - \varepsilon_b} |\bar{\psi}_{ij}^b\rangle = \frac{1}{2} \sum_{cjab} \frac{\langle ab | \hat{o} | c_j \rangle}{\varepsilon_c - \varepsilon_a + \varepsilon_j - \varepsilon_b} |\bar{\psi}_{ij}^b\rangle$$

-65-

In the following, we will sometimes use the notation,

$$\begin{aligned}\Delta^{(k)}(i_1, \dots, i_n; a_1, \dots, a_n) &= (\omega_{i_1, \dots, i_n}^{a_1, \dots, a_n})^{-k} \\ &= \left[ \sum_{j=1}^n (\varepsilon_{ij} - \varepsilon_{aj}) \right]^{-k}. \quad (36)\end{aligned}$$

Then, for example,

$$\begin{aligned}k_0^{(2)} &= \sum_{c,a} \langle c | \hat{q} | a \rangle \langle a | \hat{q} | c \rangle \Delta^{(1)}(c; a) \\ &\quad + \frac{1}{4} \sum_{c,j;a,b} \langle c j | \hat{q} | a b \rangle \langle a b | \hat{q} | c j \rangle \\ &\quad \times \Delta^{(1)}(c j; a, b) \quad (37)\end{aligned}$$

$$\begin{aligned}|H_0^{(1)}\rangle &= \sum_{c,a} \langle a | \hat{q} | c \rangle \Delta^{(1)}(c; a) |\bar{\psi}_c^a\rangle \\ &\quad + \frac{1}{4} \sum_{c,j;a,b} \langle a b | \hat{q} | c j \rangle \Delta^{(1)}(c j; a, b) |\bar{\psi}_{c j}^{a b}\rangle \quad (38)\end{aligned}$$

Note that in the H-F case ( $\hat{q}=0$ ), there is no contribution from 1p-1h excitations to  $|H_0^{(1)}\rangle$  and  $k_0^{(2)}$ . 2p-1h excitations appear already in MBPT(2) energy and MBPT(1) wave function (not a surprise for H with 2-body interaction). The question is how far do we have to go to see 3p-3h, 4p-4h, etc. excitations.

5. Third-order  $\overline{\text{MBPT}}$  correction to the energy

$$\Delta E_0^{(3)} = k_0^{(3)} = \langle \bar{\psi}_0 | W R^{(0)} W R^{(0)} W | \bar{\psi}_0 \rangle$$

$$W = W_1 + W_2 = Q_N + V_N. \quad (139)$$

There are the following groups of terms:

$$k_0^{(3)}(A) = \langle \bar{\psi}_0 | V_N R^{(0)} V_N R^{(0)} V_N | \bar{\psi}_0 \rangle, \quad (140)$$

$$k_0^{(3)}(B) = \langle \bar{\psi}_0 | V_N R^{(0)} V_N R^{(0)} Q_N | \bar{\psi}_0 \rangle$$

$$+ \langle \bar{\psi}_0 | V_N R^{(0)} Q_N R^{(0)} V_N | \bar{\psi}_0 \rangle$$

$$+ \langle \bar{\psi}_0 | Q_N R^{(0)} V_N R^{(0)} V_N | \bar{\psi}_0 \rangle, \quad (141)$$

$$k_0^{(3)}(C) = \langle \bar{\psi}_0 | Q_N R^{(0)} Q_N R^{(0)} V_N | \bar{\psi}_0 \rangle$$

$$+ \langle \bar{\psi}_0 | Q_N R^{(0)} V_N R^{(0)} Q_N | \bar{\psi}_0 \rangle$$

$$+ \langle \bar{\psi}_0 | V_N R^{(0)} Q_N R^{(0)} Q_N | \bar{\psi}_0 \rangle, \quad (142)$$


$$k_0^{(3)}(D) = \langle \bar{\psi}_0 | Q_N R^{(0)} Q_N R^{(0)} Q_N | \bar{\psi}_0 \rangle. \quad (143)$$

We begin with the A term (the only term in the Hartree-Fock case;  $Q_N = 0$ ).


$$L_0^{(3)}(A) = \langle \emptyset | V_N R^{(2)} V_N R^{(2)} V_N | \emptyset \rangle. \quad (144)$$

We have to cross all non-equivalent resulting diagrams, with no external lines and no dangerous denominators, from three  $V_N$  vertices (remembering about the denominator convention):

Nonoriented Hugenholtz skeletons:




There are 4 lines at I. If none of the lines of I goes to II, all lines of I go to III and II is left not connected. Thus, at least 1 line of I has to be connected with II. If the remaining 3 lines of I are connected with III, we get



we cannot connect all these lines.


Thus, at least 2 lines of I have to be connected with II. If  $\geq 3$  lines of I are connected

with  $\text{II}$ , we get



, from which we cannot

get a diagram with no external lines (remember,  
 $N$  is in the  $N$ -product form), thus, exactly  
2 lines must connect  $\text{I}$  and  $\text{II}$   
and exactly 2 lines meet connect  $\text{I}$  and  $\text{II}$ .  
We end up with only one scattering skeleton;




### Oriented Hugenholtz diagrams.

By introducing arrows (red arrow), we obtain:


(red is the determining

hh (hole-hole)




(i)

pp (particle-particle)






(ii)

ph-(particle-hole)




(iii)

We obtain



denominators from reduced  
resolvents



Hugenholtz

Brondum

We get the following result:

$$k_0^{(3)}(A) = k_0^{(3)}(hh) + k_0^{(3)}(pp) + k_0^{(3)}(ph),$$

(145)

where

$$k_0^{(3)}(hh) = \frac{1}{8} \sum_{ab,ijkl} \langle ab|\hat{v}|kl\rangle_A \langle kl|\hat{v}|ij\rangle_A \\ \times \langle ijl|\hat{v}|ab\rangle_A \\ \times \Delta^{(1)}(ij; \alpha, b) \Delta^{(1)}(kl; \alpha, b), \quad (146)$$





$$k_0^{(3)}(pp) = \frac{1}{8} \sum_{abcd, ij} \langle ijl|\hat{v}|ab\rangle_A \langle ab|\hat{v}|cd\rangle_A \\ \times \langle cd|\hat{v}|ij\rangle_A \\ \times \Delta^{(1)}(ij; \alpha, b) \Delta^{(1)}(ij; c, d), \quad (147)$$

$$k_0^{(3)}(ph) = - \sum_{abc, ijk} \langle bc|\hat{v}|kj\rangle_A \langle ja|\hat{v}|ib\rangle_A \\ \times \langle ik|\hat{v}|ca\rangle_A \\ \times \Delta^{(1)}(i, k; \alpha, c) \Delta^{(1)}(j, k; b, c), \quad (148)$$




Other terms contributing to  $K_0^{(3)}$  are:

$$K_0^{(3)}(B) = \langle \hat{\phi}_0 | Q_N R^{(0)} V_N R^{(0)} V_N | \hat{\phi} \rangle \\ + \langle \hat{\phi}_0 | V_N R^{(0)} Q_N R^{(0)} V_N | \hat{\phi} \rangle \\ + \langle \hat{\phi}_0 | V_N R^{(0)} V_N R^{(0)} Q_N | \hat{\phi} \rangle; \quad (149)$$



Hugenholtz diagrams:



$$K_0^{(3)}(C) = \langle \hat{\phi}_0 | Q_N R^{(0)} Q_N R^{(0)} V_N | \hat{\phi} \rangle \\ + \langle \hat{\phi}_0 | Q_N R^{(0)} V_N R^{(0)} Q_N | \hat{\phi} \rangle \\ + \langle \hat{\phi}_0 | V_N R^{(0)} Q_N R^{(0)} Q_N | \hat{\phi} \rangle; \quad (150)$$




$$K_0^{(3)}(D) = \langle \phi_0 | Q_N R^{(2)} Q_N R^{(2)} Q_N | \phi_0 \rangle; \quad (151)$$




Please note that all of these diagrams containing at least one  $Q_N$  vertex can be joined together.

Diagrams in the B group can be all obtained from



by allowing the  $Q_N$  and  $V_N$  vertices to be permuted.

Similar applies to diagrams in the C group (can all be obtained from



and in the D group (the




diagram can be obtained from




The diagrams that can be transformed into one another by topological transformations that do not break lines, but which do not preserve the order of vertices along the horizontal time axis, are referred to as the TIME VERSIONS of the same diagram.

We distinguish between:

- the versions of the first kind; vertices are permuted without changing the particle-hole character of any fermion line.



or



- time versions of the second kind:

vertices are permuted along the time axis  
and at least one line changes its p-h character.

Diagrams in each of the three groups B-D  
are in this category.

Time versions of the first kind, are very  
important for proving the linked cluster  
theorem.

In terms of physics,  $k^{(3)}$  does not bring  
information about higher than 2p-2h  
excitations. For example,  $k_6^{(3)}(A)$ ,  
which survives any type of single-particle  
basis, describes the 3rd-order contribution  
to 2p-2h excitations, since the only reduced  
residues involved are the two-body  $R_2^{(3)}$   
components. This can be easily understood  
if we realise that the two-body interaction  $V_N$   
cannot couple  $|2\rangle$  to higher than 2p-2h  
excitations.

$m, n \text{ must be } 2 \Rightarrow \langle \bar{2} | V_N R^{(0)}_{\bar{m}} V_N R^{(0)}_{\bar{n}} V_N | 2 \rangle$ .

We need to go to higher orders to see 3p-3h  
and other higher-order terms.

The remaining pages are taken directly from the lecture notes for CEM 993 class on “Algebraic and Diagrammatic Methods for Many-Fermion Systems,” taught by Piotr Piecuch at Michigan State University. The page numbers are consecutive, but they do not continue from the last page number in the preceding lecture notes prepared for the Workshop of the *Espace de Structure et de Réactions Nucléaires Théorique* on “Many-Body Perturbation Theories in Modern Quantum Chemistry and Nuclear Physics,” March 26-30, 2018, CEA Saclay, Gif-sur-Yvette, France.

Forth-order MBPT energy contributions:

$$k_0^{(4)} = \langle \emptyset | W R^{(0)} W R^{(0)} W R^{(0)} W | \emptyset \rangle - \langle \emptyset | W R^{(0)} W | \emptyset \rangle \langle \emptyset | W R^{(0)} W | \emptyset \rangle,$$

where  $W = V_N + Q_N$ .

Let us look at the purely  $V_N$  terms:

$$\langle \emptyset | V_N (R^{(0)} V_N)^3 | \emptyset \rangle \quad (\text{principal term})$$

$$- \underbrace{\langle \emptyset | V_N R^{(0)} V_N | \emptyset \rangle}_{k_0^{(2)}} \langle \emptyset | V_N R^{(0)} V_N | \emptyset \rangle \quad (\text{renorm. term})$$

Principal term:

Nonoverlapping shells



(2)

•

•

•


①

← we call these skeletons by combinatorics;  
0, 1, 2, 3, 4 lines  
between ① and ②

V2

-DSG-


I



no lines  
between ①  
and ②


[4 between ① &  
④]  
III  
 $4+0$

II




[3 between  
① & ④ and 1  
between ① and ③]  
III  
 $3+1$

III




$2+2$

IV




$1+3$

V



$0+4$


-557-

VII


1 line  
between  
① and ②

3 between  
③ and ④  
[3+0]

2+1


VIII

1+2



IX

0+3



14.



2 lines  
between  
node 1  
and 2



3 lines between  
node 1 and 3




longer  
clenomach.  
4 lines  
between node 1  
and 2

Thus, we get the following diagrams in the principal term:


## CONNECTED DIAGRAMS :

Intermediate Quotients:


II



III




IV




6.

-560-


Pterostichus  
fugax




(VI)



(VII)




(VIII)




(IX)

~~Intermediate~~ doubles:


(X)



(X)




(XII)




Intermediate singles:

(XIII)




(XIV)




DISCONNECTED D-MS in the principal form:

[I]




▽




Renormalization terms:

-



$$1/k_0^{(2)}$$

x




$$\uparrow \langle \psi | V_N R^{(0)} V_N | \psi \rangle$$

-563-

Let us see what happens with the disconnected terms from the forward form:


I



$$I = \frac{N}{b(a+b)b} \quad N \leftarrow \text{numerator}$$

(summed over  
the relevant  
spin orbit  
choices)

II



$$II = \frac{N}{a(a+b)b} \quad N \leftarrow \text{the same numerator}$$

$$I + II = \frac{N}{(a+b)b} \left( \frac{1}{a} + \frac{1}{b} \right) = \frac{N}{(a+b)b} \frac{(a+b)}{ab}$$

$$= \frac{N}{ab^2} = \frac{N}{ab^2} \times \frac{a}{a} \times \frac{b^2}{b^2}$$

As we can see, the disconnected terms from the principal term cancel the renormalization terms:

$$\begin{aligned}
 K_0^{(4)} &= \langle \bar{\phi}_0 | W(R^0 W)^3 | \bar{\phi}_0 \rangle - \langle \bar{\phi}_0 | W R^0 W | \bar{\phi}_0 \rangle \\
 &\quad \times \langle \bar{\phi}_0 | W R^0 W | \bar{\phi}_0 \rangle \\
 &= \langle \bar{\phi}_0 | \{W(R^0 W)^3\}_C | \bar{\phi}_0 \rangle + \\
 &\quad + \cancel{\langle \bar{\phi}_0 | \{W(R^0 W)^3\}_{DC} | \bar{\phi}_0 \rangle}^{\text{? connected}} \\
 &\quad - \cancel{\langle \bar{\phi}_0 | W R^0 W | \bar{\phi}_0 \rangle \langle \bar{\phi}_0 | W R^0 W | \bar{\phi}_0 \rangle}^{\text{? disconnected}}
 \end{aligned}$$

$$K_0^{(4)} = \langle \bar{\phi}_0 | \{W(R^0 W)^3\}_C | \bar{\phi}_0 \rangle$$

This is an example of a cancellation that takes place in every order and which is summarized by the **linked-cluster theorem**, which states that

$$k_S^{(n+1)} = \langle \bar{\rho}_0 | \{W(R^{\otimes n} W)^n\}_C | \bar{\rho}_0 \rangle.$$

A similar cancellation occurs in the more feynman contributions. To understand this cancellation in more feynman corrections, let us look at a few lower order contributions to  $\langle \bar{\rho}_0 \rangle$ .

We have already analyzed  $\langle \bar{\rho}_0^{(1)} \rangle$ :

$$\begin{aligned} \langle \bar{\rho}_0^{(1)} \rangle &= R^{(0)} W \langle \bar{\rho}_0 \rangle = R^{(0)} W_N \langle \bar{\rho}_0 \rangle + \\ &+ R^{(0)} Q_N \langle \bar{\rho}_0 \rangle = \text{Diagram A} + \text{Diagram B}. \end{aligned}$$


Nothing interesting happens here, both contributions to the more feynman are CONNECTED,

Let us look at  $\langle \bar{\rho}_0^{(2)} \rangle$ :

$$\langle \bar{\rho}_0^{(2)} \rangle = R^{(0)} W R^{(0)} W \langle \bar{\rho}_0 \rangle \quad (\text{There are no renormalization terms}).$$


-PGC-

Let us analyse terms that originate from  $V_{N,i}$  (Hugenholtz d<sub>m</sub>s).



(in all d<sub>m</sub>s, all lines extend to the left)

Please note that we did not draw



This would be a "dangerous denominator".

Clearly, we must always have some external lines in each of the wave function diagrams (because of the leftmost  $R^{(0)}$ ).

As we can see,

$$\begin{aligned}
 |\Psi^{(2)}\rangle &= |\Psi^{(2)}(Q)\rangle + |\Psi^{(2)}(T)\rangle \\
 &+ |\Psi^{(2)}(D)\rangle + |\Psi^{(2)}(S)\rangle
 \end{aligned}$$

↓ quadruples  $|\Psi^{(2)}_{\text{quad}}\rangle$   
 (disconnected)

triples  $|\Psi^{(2)}_{\text{trip}}\rangle$   
 singlets  $|\Psi^{(2)}_{\text{sing}}\rangle$

doubles  $|\Psi^{(2)}_{\text{dbl}}\rangle$

Triples and quadruples contribute for the first time, in the second-order MBPT wave function. Singlets contribute for the first time in  $|\Psi^{(2)}\rangle$  if H-F orbitals are used.

Diagrams contributing to  $|\Psi^{(2)}\rangle$  are of the two types: connected (slangs (ii)-(viii)) and

disconnected (from (i)). Thus, if there is a cancellation of subgraphs in the wave function, the cancellation must involve some other subgraphs than just disconnected.

Well, let us look at the 3rd order:

$$\langle \bar{q} q \rangle^{(3)} = R^{(0)} W R^{(0)} W R^{(0)} W (\bar{q} q) - \langle W R^{(0)} W \rangle R^{(0)2} W (\bar{q} q)$$

Again, let us focus on the contributions originating from  $V_N$  terms; we will draw skeletons only:

$$\underline{(R^{(0)} V_N)^3 (\bar{q} q) \text{ TERM}} \quad (\text{principal term})$$












Diagram with 3 connected parts

I   II   III





-PG-

Diagrams with 2 connected parts (I & II correct)





570


(II and III connected)









Diagrams with the connected part  
(connected oligograms)



-571



(time versions of the 2nd knot)



(time versions  
of the second  
knot)


etc.

Please notice the presence of two diagrams,  
which contain the closed (vacuum) <sup>most lines</sup>

disconnected



These two diagrams  
are two time versions of  
the first kind obtained from



(The third time version: is excluded, since  
it would lead to a dangerous denominator.


The above two diagrams are examples of the  
UNLINKED diagrams: In general, a disconnected

diagram that has at least one disconnected vacuum  
component is called UNLINKED.

LINKED Diagrams have no disconnected recessed parts.

We have the following classification of Diagrams:

CONNECTED





LINKED

DISCONNECTED



LINKED



CONNECTED

LINKED



DISCONNECTED

UNLINKED



Any centred diagram is, by definition, disconnected.

However, a forked diagram can be connected or disconnected.

We can write (returning to the  $|W_0^{(3)}\rangle$  case):

$$|W_0^{(3)}\rangle = \{ (R^{(0)} W)^3 \}_{C} |\bar{\psi}_0\rangle \quad \text{LINKED}$$

$$+ \{ (R^{(0)} W)^3 \}_{DC, L} |\bar{\psi}_0\rangle \quad \text{principal term}$$

disconnected, linked

$$+ \{ (R^{(0)} W)^3 \}_{UL} |\bar{\psi}_0\rangle \quad \text{renormalized}$$



$$- \langle \bar{\psi}_0 | W R^{(0)} W | \bar{\psi}_0 \rangle R^{(0)2} W |\bar{\psi}_0\rangle \quad \text{renorm. term}$$

The  $\{ (R^{(0)} W)^3 \}_{UL} |\bar{\psi}_0\rangle$  part is represented by two time versions of the first kind contact-like 

Mechanism part:

-575

$a \{$   
 $b \}$



$$(I) = \frac{N}{b(a+b)b}$$

$$(II) = \frac{N}{b(a+b)a}$$

↗ ( summed over the relevant spin-orbital labels ).

[ $N$  is the numerator, i.e. the product of the  $v$  matrix elements and  $Y^+$  operators corresponding to external lines, sign, and weight factors]

$$\begin{aligned} \{(RCW)^3\}_{ab}(\tilde{\rho}_0) &= \frac{N}{b(a+b)b} + \frac{N}{b(a+b)a} \\ &= \frac{N}{b(a+b)} \left( \frac{1}{b} + \frac{1}{a} \right) = \frac{N}{b(a+b)} \cancel{\frac{ab}{ab}} ab \\ &= \frac{N}{ab^2} = a \{ \text{Diagram (I)} \} b \times b \{ \text{Diagram (II)} \} \end{aligned}$$

-576

$$= \langle \hat{\phi}_0 | W R^{(0)} W | \hat{\phi}_0 \rangle \cancel{R^{(0)} W} | \hat{\phi}_0 \rangle.$$

Thus, the unlinked part of  $| \hat{\phi}_0^{(3)} \rangle$  cancels the renormalization term and we obtain:

$$| \hat{\phi}_0^{(3)} \rangle = \{ (R^{(0)} W)^3 \}_C | \hat{\phi}_0 \rangle$$

$$+ \{ (R^{(0)} W)^3 \}_{DC\cancel{L}} | \hat{\phi}_0 \rangle = \{ (R^{(0)} W)^3 \}_{\cancel{L}} | \hat{\phi}_0 \rangle$$

↑  
linked

---

A very similar cancellation of unlinked principal and renormalization terms takes place in every order,

$$| \hat{\phi}_0^{(n)} \rangle = \{ (R^{(0)} W)^n \}_{\cancel{L}} | \hat{\phi}_0 \rangle,$$

Only linked diagrams

For example, in the 4th order, the unlinked principal terms are:


-577-

$$\{(R^{\odot} W)^4\}|\Phi_0\rangle \Rightarrow$$

UL




(i)



+ tube versions  
of the  
1st lens

+


(ii)



+ tube versions  
of the 2nd  
lens


+

(iii)



+ tube versions of  
the 3rd lens

(iv)




+ tube versions of  
the 4th lens

-578-

5

(v)

+








+ fine version of  
the (A) knot

-579-

Lotus by group (iii):

a)



||

$$= \frac{N}{b(a+b)bc} + \frac{N}{b(a+b)(a+c)c} + \frac{N}{b(a+b)(a+c)a}$$

$$+ \frac{N}{bc(a+c)c} + \frac{N}{bc(a+c)a} =$$

-80-

$$= \frac{N}{b(a+b)bc} + \frac{N}{b(a+b)(a+c)} \left( \frac{1}{c} + \frac{1}{a} \right)$$

$$+ \frac{N}{bc(a+c)} \left( \frac{1}{c} + \frac{1}{a} \right) =$$

$$= \frac{N}{b(a+b)bc} + \frac{N}{b(a+b)(a+c)} \cancel{\frac{(a+c)}{ac}}$$

$$+ \frac{N}{bc(a+c)} \cancel{\frac{(a+c)}{ac}} =$$

$$= \frac{N}{b(a+b)bc} \quad \leftarrow \quad + \frac{N}{b(a+b)ac} + \frac{N}{abc^2}$$

$$= \frac{N}{b(a+b)c} \left( \frac{1}{b} + \frac{1}{a} \right) + \frac{N}{abc^2}$$

$$= \frac{N}{b(a+b)c} \cancel{\frac{(a+b)}{ab}} + \frac{N}{abc^2}$$

$$= \frac{N}{ab^2c} + \frac{N}{abc^2} =$$

→ 58 (-)

$$= \text{(a)} \times \left( \begin{array}{c} \text{(b)} \\ \text{(c)} \end{array} \right) + \text{(b)} \times \text{(c)}$$

By continuing a similar analysis for the remaining UQ domes, we obtain:

$$\left\{ (R^{\infty} W)^3 \right\}_{UQ} (\phi_0) =$$

→ 882-

$$= \langle WR^{(0)}W \rangle [R^{(0)2}WR^{(0)}W |\bar{\phi}\rangle]$$

$$+ R^{(0)}W R^{(0)2}W |\bar{\phi}\rangle]$$

$$+ \langle WR^{(0)}WR^{(0)}W \rangle R^{(0)2}W |\bar{\phi}\rangle$$

= renormalization forms in

$$\langle \bar{\phi}^{(4)} \rangle$$

Thus, again,

$$\langle \bar{\phi}^{(4)} \rangle = \{ (R^{(0)}W)^3 \}_W |\bar{\phi}\rangle$$

Please note that in order for the above cancellations to take place, we must assume that all labels in the diagrams correspond to unrenormalized quantities. Indeed,

$$\{ (R^{(0)}W)^3 \}_W |\bar{\phi}\rangle = \sum_{i,j,\dots} \left( \text{diagram } i + \text{diagram } j \right)$$

$$= \sum_{i,j,\dots} \text{diagram } i \times \text{diagram } j = \left( \sum_i \text{diagram } i \right) \sum_j \text{diagram } j$$

(8) / (R^{(0)2}W)

coll's      coll's

coll's      coll's

83-

The renormalization form -  $\langle W R^{\text{co}} W \rangle R^{\text{co}} W | \frac{\partial}{\partial} \rangle$ .  
These summations are unrestricted, since we replaced the original formula for  $R^{\text{co}}$ ,

$$R^{\text{co}} = \sum_{n=1}^N R_n^{(0)}$$

$$R_n^{(0)} = \sum_{\substack{\text{restricted} \\ \{i_1, i_2, \dots, i_n \\ i_1 \neq i_2 \neq \dots \neq i_n \\ \alpha_1 \neq \alpha_2 \neq \dots \neq \alpha_n}}} \frac{\langle \Phi_{i_1-i_n}^{a_1-a_n} \rangle \langle \Phi_{i_1-i_n}^{a_1-a_n} |}{-\delta_{i_1-i_n}^{a_1-a_n}}$$

by

$$R_n^{(0)} = \left(\frac{1}{n!}\right)^2 \sum_{\substack{\text{unrestricted} \\ \{i_1-i_n \\ a_1-a_n \\ i_1 \neq i_2 \neq \dots \neq i_n \\ \alpha_1 \neq \alpha_2 \neq \dots \neq \alpha_n\}}} \frac{\langle \Phi_{i_1-i_n}^{a_1-a_n} \rangle \langle \Phi_{i_1-i_n}^{a_1-a_n} |}{-\delta_{i_1-i_n}^{a_1-a_n}}$$

Clearly, all terms with  $i_1 = i_2$  or  $i_1 = i_3$ , etc., in  $R^{\text{co}}$  vanish, but when we form more complicated quantities using diagrams they will contribute (although, mutually cancel out in the final diagrams).

Suppose, we used

$$R_n^{(0)} = \left(\frac{1}{n!}\right)^2 \sum_{\substack{i_1 \neq i_2 \neq \dots \neq i_n \\ \alpha_1 \neq \alpha_2 \neq \dots \neq \alpha_n}} \frac{\langle \Phi_{i_1-i_n}^{a_1-a_n} \rangle \langle \Phi_{i_1-i_n}^{a_1-a_n} |}{-\delta_{i_1-i_n}^{a_1-a_n}}$$

In the first order, we would get

- 584 -

$$\{ (R^{\otimes} W)^3 \}_{U \subset \{i\}} |\tilde{\psi}_0\rangle = \sum_{i \neq j} ( \text{Diagram with } i \text{ and } j \text{ connected by a horizontal line} ) + ( \text{Diagram with } i \text{ and } j \text{ connected by a vertical line} )$$

$$= \sum_{i \neq j} ( \text{Diagram with } i \text{ and } j \text{ connected by a horizontal line} ) + ( \text{Diagram with } i \text{ and } j \text{ connected by a vertical line} )$$

$$- \sum_{i \dots} ( \text{Diagram with } i \text{ and } j \text{ connected by a horizontal line} ) + ( \text{Diagram with } i \text{ and } j \text{ connected by a vertical line} ) \quad (\leftarrow EPV \text{ terms})$$

$$= \langle W R^{\otimes} W \rangle R^{\otimes 2} W |\tilde{\psi}_0\rangle -$$

$$- \sum_{i \dots} ( \text{Diagram with } i \text{ and } j \text{ connected by a horizontal line} ) \times ( \text{Diagram with } i \text{ and } j \text{ connected by a vertical line} ), \text{ so that}$$

$$|\tilde{\psi}_0\rangle = \{ (R^{\otimes} W)^3 \}_{\{i\}} |\tilde{\psi}_0\rangle + \{ (R^{\otimes} W)^3 \}_{U \subset \{i \neq j\}} |\tilde{\psi}_0\rangle$$

$$- \langle W R^{\otimes} W \rangle R^{\otimes 2} W |\tilde{\psi}_0\rangle$$

↓ unlabeled EPV

$$= \{ (R^{\otimes} W)^3 \}_{U \subset \{i \neq j\}} |\tilde{\psi}_0\rangle = \sum_{i \dots} ( \text{Diagram with } i \text{ and } j \text{ connected by a horizontal line} ) + ( \text{Diagram with } i \text{ and } j \text{ connected by a vertical line} )$$

-88-

The latter term is unlinked and is left uncanceled, since we restricted the summations.


By adding the linked  $i = j$  terms to the sum

$$\{(R^{\otimes} W)^3\}_{(i=j)} | \Psi \rangle ,$$

we can immediately eliminate the unlinked


$\sum_{i=1}^n$   


term. Instead, by

 adding  $\sum \{(R^{\otimes} W)^3\}_{(i=j)} | \Psi \rangle$  to  $| \Psi^{(3)} \rangle$ ,


we obtain:

$$| \Psi^{(3)} \rangle = \underbrace{\{(R^{\otimes} W)^3\}_{(i=j)} | \Psi \rangle}_{\text{null}} - \{(R^{\otimes} W)^3\}_{(i=j)} | \Psi \rangle$$

$- \sum_i$  





 will cancel each other





86-

$\{(R^{\text{eff}})^3\}_{(i=j)}[\ell_0]$  contains




These terms cancel unlinked renormalization  $i=j$  forms:



(Indeed, (IIA))

(Brenda)



$$-\frac{1}{4} \sum_{cd} \langle \text{eff}|\rho|l;i\rangle \langle \text{coll}|v|k;j\rangle$$


$$\times \langle k|i|\delta|cd\rangle_A$$

$$\times \Delta^{(1)}(i,l;\text{eff})$$

$$\times \Delta^{(1)}(i,k,l;c,d,\text{eff})$$

$$\times \Delta^{(1)}(e,l;\text{eff})$$

(IIA)



$$\frac{1}{4} \sum_{\text{config.}} \langle \text{eff.} | l_i \rangle$$

$$x \langle \text{cd} | \beta | k_i \rangle \langle \text{cd} | \beta | d \rangle$$

$$x \Delta^D(i,l;ef) \Delta^D(i,k,l;e,d,ef) \Delta^D(i,l;ef)$$

Thus,

$$(IA) + (IIA) = 0.$$

Similarly,  $(IB) + (IIB) = 0$ .



Thus, after adding and subtracting the terms

$i=j$  terms,

$$|I_0^{(3)}\rangle = \{(R^{\infty}W)^3\} |B\rangle.$$

This example illustrates the need for consistency. The so-called exclusion principle prohibits (EPV) subgraphs, in which a given symbol state is occupied more than once (we have two identically labeled hole or particle lines).

In general, the EPV terms cancel out.  
Schematically,



EPV I - m

another EPV I - m.

(number of legs changes before here)

Clearly, the above diagrams cancel out. They do not have to vanish, but once all of the ~~gluograms~~<sup>in the principal renormalization</sup>, EPV, are considered, EPV terms mutually cancel out.

In some cases, EPV diagrams that cancel out are in the principal form. However, there are cases where one of the two DV diagrams that cancel out is linked and another unlinked.

In the above example, DV diagram (I A) is linked, EPV diagram (II A) is unlinked. Yet they cancel out. If we did not allow the EPV terms, the above cancellations of unlinked principal and renormalization terms would not be complete. Thus, the linked outer theorem will have a form:

$$|\Psi^{(n)}\rangle = \sum_{\substack{\text{L, including EPV} \\ \text{L, excluding EPV}}} \{(R^{\odot n})^L\}$$

(B)

The vanished terms involving EPV diagrams cancel out. ~~Other~~ statements, such as:

$$|\Psi^{(n)}\rangle = \sum_{\text{no EPV}} \{(R^{\odot n})^L\}$$

(C)


are not true.

## 67. Factorization Lemma

and the connected cluster theorem

Before proving the linked diagram (cluster) theorem, we have to prove the so-called Factorization Lemma (following the work of Franks and Mills). This lemma allows to factorize the disconnected, but linked diagrams, a construction appearing in the proof of the linked cluster theorem. These diagrams are obtained in the ~~Let us illustrate the Factorization Lemma by a few examples.~~


Suppose we have a



proof of the linked cluster theorem by removing the leftmost cluster vertex from the ~~nonseparating part~~ of the unlabeled diagrams having precisely one nonseparating part).

Let us illustrate the Factorization Lemma by a few examples:


- disconnected linked diagrams having nonequivalent connected components:




+ all home versions  
of the first kind

nonequivalent


-59)



+



+




$$\frac{N}{(a+b)bc} + \frac{N}{(a+b)(a+c)c} + \frac{N}{(a+b)(a+c)a}$$

$$= \frac{N}{(a+b)bc} + \frac{N}{(a+b)(a+c)} \left( \frac{1}{a} + \frac{1}{c} \right)$$


$$= \frac{N}{(a+b)bc} + \frac{N}{(a+b)ac} = \frac{N}{(a+b)c} \left( \frac{1}{b} + \frac{1}{a} \right)$$

$$= \frac{N}{abc}$$


Denominator in the  
first term




$\times$



- ④ Disconnected linked diagrams having equivalent connected components:



This diagram does not ~~have~~ have nonequivalent line versions, but we can obtain the same ~~diagram~~ combination by inferring the above diagram twice, in two equivalent line versions (all graphically labeled as per), and by dividing the result by 2.




$$\begin{aligned} & \frac{1}{2} \left( \frac{N}{(a+b)b} + \frac{N}{(a+b)a} \right) = \frac{1}{2} \frac{N}{(a+b)} \left( \frac{1}{b} + \frac{1}{a} \right) \\ & = \frac{1}{2} \frac{N}{ab} \end{aligned}$$

over, the sum on the left proven

Thus, when we have two equivalent parts, we obtain a similar factorization as in the earlier example, but we also get a factor of  $\frac{1}{2}$  associated with the part that we have two equivalent parts. But it's consistent with our rules for topological factors, since disconnected equivalent parts ~~can be permuted~~ among themselves when factored and counted as independent components.


The above example was ~~somewhat simple~~, very simple, both connected parts were simple vectors. Let us try something more complicated:

- disconnected linked diagrams having equivalent connected components (a more complicated example):




+ all fine versions  
nonequivalent

(we must assume that both connected pieces have the same orientation of lines; say left, if one is left and the other right, both parts are right)



If we enlarged the above three diagrams, we would not achieve saturation; however, we can double the number of diagrams by combining all four varieties of the 1st kind and doubling by a factor of 2.



$$(I = \bar{V}; \quad \bar{II} = \bar{V}; \quad \bar{III} = \bar{V}).$$

$$\begin{aligned}
 &= \frac{1}{2} \left\{ \frac{N}{(a+c)(b+d)} a + \frac{N}{(a+c)(b+c)(b+d)} \cancel{a} + \frac{N}{(a+c)(b+c)(b+d)} b \right. \\
 &\quad + \frac{N}{(a+c)(a+d)(b+d)} \cancel{a} + \frac{N}{(a+c)(a+d)(b+d)} b \\
 &\quad \left. + \frac{N}{(a+c)(a+d)a b} \right\} =
 \end{aligned}$$

-595-




$$= \frac{1}{2} \left\{ \frac{N}{(a+c)(b+d)c\bar{d}} + \frac{N}{(a+c)(b+d)\bar{b}\bar{d}} \right\}$$

$$+ \frac{N}{(a+c)(a+d)\bar{b}\bar{d}} + \frac{N}{(a+c)(a+d)a\bar{b}} \right\}$$

$$= \frac{1}{2} \left\{ \frac{N}{(a+c)\bar{d}\bar{b}\bar{c}} + \frac{N}{(a+c)b\bar{a}\bar{d}} \right\}$$


$$= \frac{1}{2} \frac{N}{(a+c)b\bar{d}} \left( \frac{1}{c} + \frac{1}{a} \right)^{\frac{a+c}{ab}}$$

$$= \frac{1}{2} \frac{N}{abc\bar{d}}$$



At home: examine books with more than 2 equivalent paths.

Show that



$$= \frac{1}{6} (3!)^3$$

Let us generalize the above result:


Consider all possible fine versions of the first kind for a LINKED DISCONNECTED diagram consisting of two ports: A and B. These two ports are not necessarily connected, so the ~~rule~~ applies to ALL LINKED DISCONNECTED diagrams.

We designate the set of energy denominators for port A alone by

$\Delta_\mu^A$ ,  $\mu = 1, \dots, m$ , and for

port B by:  $\Delta_\nu^B$ ,  $\nu = 1, \dots, n$ .

The denominators are inserted along the fine axis, i.e., the rightmost denominators are  $\Delta_1^A$  and  $\Delta_1^B$ .



The denominator contribution from all possible time versions of the first kind, corresponding to all possible exten~~sions~~<sup>interactions</sup> of vertices in parts A and B relative to one another (the memory part is always Montreal for all time versions) can be written as:

$$D_{mn}^{AB} = \sum_{\{\alpha, \beta\}} \prod_{p=1}^{m+n} (\Delta_{\alpha(p)}^A + \Delta_{\beta(p)}^B)^{-1},$$

where the summation over  $\alpha, \beta$  extends over all sets of  $(m+n)$  integer pairs,

$$\Gamma_p = (\alpha(p), \beta(p)), \quad 0 \leq \alpha(p) \leq m, \\ 0 \leq \beta(p) \leq n,$$

defined as follows:

$$(i) \quad \Gamma_1 = (1, 0) \text{ or } (0, 1),$$

$$(ii) \quad \Gamma_{p+1} = (\alpha(p)+1, \beta(p)) \text{ or} \\ (\alpha(p), \beta(p)+1),$$

$$(iii) \quad \Gamma_{m+n} = (m, n).$$



We also define:

$$\Delta_0^A = \Delta_0^B = 0.$$

598

Explanation:

(i) This condition reflects the fact that the rightmost perturbation vertex is either from part A [ $\Gamma_1 = (1, 0)$  case] or from part B [ $\Gamma_1 = (0, 1)$ ].



$$\Gamma_1 = (1, 0)$$

$$\text{denominator: } \Delta_1^A + \Delta_0^B = \Delta_1^A$$




$$\Gamma_1 = (0, 1)$$

$$\text{denominator: } \Delta_n^A + \Delta_1^B = \Delta_1^B$$

~~59~~

(ii) After going through the first  $p$  vertices, the next vertex can be either in the A part or in the B part, e.g.



A (denoms:  $\Delta_1^A, \Delta_2^A$ )

B (denoms:  $\Delta_1^B, \Delta_2^B$ )

$$\Gamma_2 = (1, 1) \quad \Gamma_1 = (1, 0)$$

$$\Gamma_4 = (2, 2) \quad \Gamma_3 = (2, 1)$$

Denominators:

$$(\Delta_1^A + \Delta_0^B)^{-1} (\Delta_1^A + \Delta_1^B)^{-1} (\Delta_2^A + \Delta_1^B)^{-1}$$

$$\times (\Delta_2^A + \Delta_2^B)^{-1}$$

(iii) The last (leftmost) denominator is

$$(\Delta_m^A + \Delta_n^B)^{-1}$$

(cf., the above example),  
on (ii)

→ GDD

For the separate parts A and B, the denominators are given by the products of  $\Delta_{\mu}^A$  and  $\Delta_{\nu}^B$ , respectively,

$$D_m^A = \prod_{\mu=1}^m (\Delta_{\mu}^A)^{-1},$$

$$D_n^B = \sum_{\nu=1}^n (\Delta_{\nu}^B)^{-1}.$$

Note that

$$D_m^{AB} = D_{m0}^{AB}, \quad D_n^B = D_{0n}^{AB}.$$

Indeed:

$$\begin{aligned} D_{m0}^{AB} &= \sum_{\{\alpha, \beta\}} \prod_{p=1}^{m+0} (\Delta_{\alpha(p)}^A + \Delta_{\beta(p)}^B)^{-1} \\ &= \sum_{\{\alpha, \beta\}} \prod_{p=1}^m (\Delta_{\alpha(p)}^A + \Delta_{\beta(p)}^B)^{-1} \\ &\equiv \left( 0 \leq \beta(p) \leq n; \beta(p) = 0 \forall p \Rightarrow \Delta_{\beta(p)}^B = \Delta_0^B = 0 \right) \end{aligned}$$

$$= \sum_{\{\alpha\}} \prod_{p=1}^m [\Delta_{\alpha(p)}^A]^{-1}. \quad \text{But,}$$

since  $\beta(p) = 0$ ,

-601-

$$\Gamma_1 = (1, 0), \text{ so that } \alpha(1) = 1,$$

$$\Gamma_2 = (\alpha(1)+1, \beta(1)) = (1+1, 0) = (2, 0), \text{ so that } \alpha(2) = 2,$$

$$\Gamma_{p+1} = (\alpha(p)+1, \beta(p)) = (\alpha(p)+1, 0) = (\alpha(p+1)),$$

$$\text{so that } \alpha(p+1) = \alpha(p) + 1, \text{ which}$$

$$\text{means that } \alpha(p) = p, p = 1, \dots, m,$$

which in turn implies that

$$D_{mn}^{AB} = \prod_{p=1}^m (\Delta_p^A)^{-1} = D_m^A.$$

$$\text{Similarly for } D_n^B = D_{0n}^{AB},$$

We also define:

$$D_0^A = D_0^B = D_{00}^{AB} = 1.$$

Factorization lemma states that

$$D_{mn}^{AB} = D_m^A D_n^B$$

sum of the  
denominators from all the ratios

-602-

Proof: Mathematical induction;

$m=0$  or  $n=0$ :

$$D_{m0}^{AB} = D_m^A = D_m^A D_0^B$$

$$D_{0n}^{AB} = D_n^B = D_0^A D_n^B$$

Check on  $D_{II}^{AB}$  (e.g.) he

$\Gamma_2 = (1,1) \quad \Gamma_1 = (0,1)$        $\Gamma_2 = (1,1) \quad \Gamma_1 = (1,0)$

$$D_{II}^{AB} = \sum_{\{\alpha, \beta\}} \prod_{p=1}^2 \left[ \Delta_{\alpha(p)}^A + \Delta_{\beta(p)}^B \right]^{-1}$$

$$= \sum_{\{\alpha, \beta\}} \left[ \Delta_{\alpha(1)}^A + \Delta_{\beta(1)}^B \right]^{-1} \left[ \Delta_{\alpha(2)}^A + \Delta_{\beta(2)}^B \right]^{-1}$$

$$= (\Delta_1^A + \Delta_0^B)^{-1} (\Delta_1^A + \Delta_1^B)^{-1}$$

$$+ (\Delta_0^A + \Delta_1^B)^{-1} (\Delta_1^A + \Delta_1^B)^{-1}$$

$$= [(\Delta_1^A)^{-1} + (\Delta_1^B)^{-1}] (\Delta_1^A + \Delta_1^B)^{-1} =$$

-603-


$$= (\Delta_1^A)^{-1} (\Delta_1^B)^{-1} \cancel{(\Delta_1^A + \Delta_1^B)} (\cancel{\Delta_1^A + \Delta_1^B})^{-1}$$

$$= (\Delta_1^A)^{-1} (\Delta_1^B)^{-1} = D_1^A D_1^B .$$

(Induction step:


Suppose the lemma holds for  $M = m-1, N = n$  and  $M = m$  and  $N = n-1$ ;  $m, n \geq 1$ ,  
Let us consider the  $(m, n)$  case.

All terms in  $D^{AB}$  can be shifted onto two distant classes depending on whether the leftmost intersection occurs in  $A$  or  $B$ ; schematically,



$$\begin{aligned} &= (m, n) \\ &= (m+n, m+n-1) \end{aligned}$$

(I)



$$\begin{aligned} &= (m, n) \\ &= (m+n, m+n-1) \end{aligned}$$

(II)

- 604.

The left denominator (the leftmost one) is  
(see (iii)):

$$(\Delta_m^A + \Delta_n^B)^{-1} \quad (\Gamma_{m+n} = (m, n) \Rightarrow$$
$$\Rightarrow \alpha(m+n) = m, \beta(m+n) = n).$$

The remaining part of  $D_{mn}^{AB}$  is either  
 $D_{m-1, n}^{AB}$  (case I) or  $D_{m, n-1}^{AB}$  (case II),  
since after pulling out  $(\Delta_m^A + \Delta_n^B)^{-1}$   
out the remaining ~~denominator~~ are identical to those  
obtained for the diagrams obtained by deleting  
the leftmost vertex. Thus,

$$D_{mn}^{AB} = (\Delta_m^A + \Delta_n^B)^{-1} [ D_{m-1, n}^{AB} + D_{m, n-1}^{AB} ]$$

From the induction assumption,

$$D_{m-1, n}^{AB} = D_{m-1}^A D_n^B,$$

- 60

$$D_{m,n-1}^{AB} = D_m^A D_{n-1}^B.$$

Now,

$$D_m^A = D_{m-1}^A (\Delta_m^A)^{-1},$$

$$D_n^B = D_{n-1}^B (\Delta_n^B)^{-1}$$

Thus,

$$\begin{aligned} D_{mn}^{AB} &= (\Delta_m^A + \Delta_n^B)^{-1} [D_{m-1}^A D_n^B + D_{m,n-1}^{AB}] \\ &= (\Delta_m^A + \Delta_n^B)^{-1} [ \Delta_m^A D_m^A D_n^B + \Delta_n^B D_m^A D_n^B ] \\ &= (\Delta_m^A + \Delta_n^B)^{-1} (\Delta_m^A + \Delta_n^B) D_m^A D_n^B \\ &= D_m^A D_n^B. \end{aligned}$$

This completes the proof.

## LINKED CLUSTER THEOREM :

We have already postulated the linked cluster theorem (based on examples) in the following way:

$$|\Psi_0^{(n)}\rangle = \{ (R^{\odot n})^n \}, |\Phi\rangle$$

Tell linked sl-ms, involving  
CPV


This implies that

$$\begin{aligned} k_0^{(n+1)} &= \langle \Phi | W | \Psi^{(n)} \rangle \\ &= \underbrace{\langle \Phi | W \{ (R^{\odot n})^n \} }_{\text{no vacuum parts}}, |\Phi\rangle \\ &= \langle \Phi | (W \{ (R^{\odot n})^n \})_{C_0} |\Phi\rangle \end{aligned}$$

connected, no external  
lines, since

W has to connect to external lines of  
 $\{ (R^{\odot n})^n \}$ ; ~~if the  $(R^{\odot n})^n$  diagram is correct,~~

~~we obviously get the  $C_0$  terms;~~  
~~if  $(R^{\odot n})^n$  is linked but~~  
~~disconnected, we have~~



and all lines of A and B must be

-607-

connected with  $W$ , producing the connected oligograms.

Thus,

$$k_0^{(n+1)} = \langle \Phi_0 | \{W(R^{(0)}H)^n\}_{C_0} | \Phi_0 \rangle.$$

In other words:

$$k_0 - \Phi_0 = \sum_{n=0}^{\infty} k_0^{(n+1)} =$$

$$= \sum_{n=0}^{\infty} \langle \Phi_0 | \{W(R^{(0)}H)^n\}_{C_0} | \Phi_0 \rangle$$

$$\langle \tilde{\Phi} \rangle = \sum_{n=0}^{\infty} \langle \tilde{\Phi}_0 | \{ (R^{(0)}H)^n \}_{C_0} | \tilde{\Phi}_0 \rangle$$

Proof of the linked cluster theorem:

We must show that

$$(k_0 + W) \langle \tilde{\Phi} \rangle = k_0 \langle \tilde{\Phi} \rangle$$

for  $\langle \tilde{\Phi} \rangle$  and  $k_0$  defined above.

-GOF

Let us calculate

$$(\alpha_0 - K_0) \langle \tilde{\psi}_0 \rangle \quad (\text{using } |\tilde{\psi}_0\rangle \text{ defined by the linked terms}):$$

$$(\alpha_0 - K_0) \langle \tilde{\psi}_0 \rangle =$$

$$= (\alpha_0 - K_0) \left[ \langle \tilde{\psi}_0 \rangle + \sum_{n=1}^{\infty} \{ (R^{(0)} W)^n \} \langle \tilde{\psi}_0 \rangle \right]$$


$$= (\alpha_0 - K_0) \sum_{n=1}^{\infty} \{ (R^{(0)} W)^n \} \langle \tilde{\psi}_0 \rangle$$

$$= (\alpha_0 - K_0) R^{(0)} \sum_{n=0}^{\infty} \{ W (R^{(0)} W)^n \}_{\text{ext}} \langle \tilde{\psi}_0 \rangle$$

Since  $\langle \tilde{\psi}_0 \rangle$  are all <sup>linked</sup> vertex terms with

external lines. There must be external lines in

$W (R^{(0)} W)^n$  since we must connect lines of  $W (R^{(0)} W)^n$



( $R^{(0)}$  has at least two lines extending to the right).

Recall that

$$(\alpha_0 - K_0) R^{\text{co}} = 1 - \langle \bar{\psi} \rangle \langle \psi \rangle.$$

Thus,

$$(\alpha_0 - K_0) |\bar{\psi}_0 \rangle = (1 - \langle \bar{\psi} \rangle \langle \psi \rangle)$$

$$\leftarrow \sum_{n=0}^{\infty} \{ W (R^{\text{co}} W)^n \}_{\text{ext}} |\bar{\psi}_0 \rangle$$

$$= \sum_{n=0}^{\infty} \{ W (R^{\text{co}} W)^n \}_{\text{ext}} |\bar{\psi}_0 \rangle$$

$$- |\bar{\psi}_0 \rangle \sum_{n=0}^{\infty} \underbrace{\{ \cancel{\bar{\psi}} \cancel{\psi} (R^{\text{co}} W)^n \}}_{\text{Lext}} |\bar{\psi} \rangle,$$

"0, since Lext & ms  
have external lines.

so that

$$(\alpha_0 - K_0) |\bar{\psi} \rangle = \sum_{n=0}^{\infty} \{ W (R^{\text{co}} W)^n \}_{\text{ext}} |\bar{\psi}_0 \rangle$$

-610-

Clearly, we can only obtain the Lext  
diagrams  $\{W(R^{\otimes n})\}_{\text{Lext}} |\bar{\psi}\rangle$

from the linked diagrams

$\{(R^{\otimes n})\}_{\text{Lext}}$  (adding W)

to the UL diagram cannot produce the linked  
diagram).

Thus,

$$(\mathcal{E}_0 - K_0) |\bar{\psi}_0\rangle = \sum_{n=0}^{\infty} \left\{ W \{ (R^{\otimes n}) \}_{\text{Lext}} \right\} |\bar{\psi}_0\rangle$$

linked

linked with  
external lines


Let us analyse what we get by applying W  
to the  $\{(R^{\otimes n})\}_{\text{Lext}}$  terms:

$$W \sum_{n=0}^{\infty} \left\{ (R^{\otimes n}) \right\}_{\text{Lext}} |\bar{\psi}_0\rangle =$$

-61-

$$= \sum_{n=0}^{\infty} \{W\}_{L} \{R^{(0)} W\}^n \}_{C_0} \langle \bar{\phi} \rangle$$

$W$  is used to close all the external lines of  $\{R^{(0)} W\}^n_L$ , as in




This must be connected, since  $A$  does not have vacuum parts

$$+ \sum_{n=0}^{\infty} \{W\}_{L} \{R^{(0)} W\}^n \}_{W^{\text{ext}}} \langle \bar{\phi} \rangle$$

$W^{\text{ext}}$  entwined with external lines

$W$  is closing the cones of one of the disconnected parts of  $\{R^{(0)} W\}^n$  (producing the vacuum term), producing enriched diagrams (having external lines)



$W^{\text{ext}}$  (enriched),  
 $B$  has external lines

$$+ \sum_{n=0}^{\infty} \left\{ W \left\{ (R^{\odot} W)^n \right\}_L \right\}_{\text{Ext}} |\tilde{\Psi}_0\rangle$$

$W$  is not closing the loops of any of the parts of  $\{(R^{\odot} W)^n\}_L$ , leaving us with bounded diagrams with some external lines.

This means that

$$(s_0 - k_0) |\tilde{\Psi}_0\rangle = W \sum_{n=0}^{\infty} \left\{ (R^{\odot} W)^n \right\}_L |\tilde{\Psi}_0\rangle$$

$$- \sum_{n=0}^{\infty} \left\{ W \left\{ R^{\odot} W \right\}^n \right\}_L |_{C_0} |\tilde{\Psi}_0\rangle$$

$$- \sum_{n=0}^{\infty} \left\{ W \left\{ (R^{\odot} W)^n \right\}_L \right\}_{\text{Ext}} |\tilde{\Psi}_0\rangle$$

Now:

$$\sum_{n=0}^{\infty} \left\{ W \left\{ (R^{\odot} W)^n \right\}_L \right\}_{C_0} |\tilde{\Psi}_0\rangle$$

$$= \sum_{n=0}^{\infty} \left\{ W (R^{\odot} W)^n \right\}_{C_0} |\tilde{\Psi}_0\rangle = (k_0 - s_0) \times |\tilde{\Psi}_0\rangle.$$


-673 -

$$\sum_{n=0}^{\infty} \{W\{(R^{\odot}H)^n\}\}_{\text{ULext}} \stackrel{(\oplus)}{>} =$$

(in reality  $n \geq 1$ )



$$\sum_{i,j,a,b}$$




+ all tame versions

$$\sim \{(R^{\odot}H)^n\}_{2 \text{ (or more)}}$$

=


$$\sum_{i,j,a,b}$$




## FACTORIZATION

LEMMA

$$\sum_{i,j,a,b}$$



=




Since we consider ALL ORDERS in

$$\sum_{n=0}^{\infty} \{W\{(R^{\odot}H)^n\}\}_{\text{ULext}} \stackrel{(\oplus)}{>},$$

the


$\{(R^{\odot}H)^n\}_{\text{ULext}}$  with  $n \geq 1$ , so that after combining

-Q4-

with  $W$ ,  contains all CONNECTED vacuum diagrams of the  $\{W\{(R^0 W)^n\}\}_{C_0}$  type. In other words, all  terms give  $(k_0 - \delta_0)$ .

Since we have all orders in

$$\sum_{n=0}^{\infty} \{W\{(R^0 W)^n\}\}_{UL_{ext}} |\bar{\Phi}_0\rangle$$

The  piece represents all linked terms with external lines, i.e.

$$\sum_{n=1}^{\infty} \{(R^0 W)^n\}_{UL} |\bar{\Phi}_0\rangle = |\bar{\Phi}_0\rangle - |\bar{\Phi}_S\rangle.$$

This implies that

$$\begin{aligned} & \sum_{n=0}^{\infty} \{W\{(R^0 W)^n\}\}_{UL_{ext}} |\bar{\Phi}_0\rangle \\ &= (k_0 - \delta_0) (|\bar{\Phi}_0\rangle - |\bar{\Phi}_S\rangle). \end{aligned}$$

-Q5-

From the above equations for

$$\sum_{n=0}^{\infty} \{W\{(R^{\infty}H)^n\}_L\}_{C_0} \langle \Psi_0 \rangle \text{ and}$$

$$\sum_{n=0}^{\infty} \{W\{(R^{\infty}H)^n\}_L\}_{ULext} \langle \Psi_0 \rangle \text{ terms, we}$$

note,

$$(x_0 - k_0) \langle \Psi_0 \rangle = W \langle \Psi_0 \rangle - (k_0 - x_0) \langle \Psi_0 \rangle$$
$$- (k_0 - x_0) (\langle \Psi_0 \rangle - \langle \Psi_0 \rangle) =$$
$$= W \langle \Psi_0 \rangle - (k_0 - x_0) \langle \Psi_0 \rangle$$

$$(x_0 - k_0) \langle \Psi_0 \rangle = W \langle \Psi_0 \rangle -$$
$$- (k_0 - x_0) \langle \Psi_0 \rangle$$

$$(k_0 + W) \langle \Psi_0 \rangle = k_0 \langle \Psi_0 \rangle,$$

---

$$\text{i.e., the } \langle \Psi_0 \rangle = \sum_{n=0}^{\infty} \{(R^{\infty}H)^n\}_L \langle \Psi_0 \rangle$$

more generally satisfies the Schrödinger equation -  
thus completes the proof!

-616-

We proved the linked cluster (diagram) theorem, which states that

$$|\psi^{(n)}\rangle = \{ (R^{\odot}W)^n \}_c |\Psi\rangle,$$

$$k_0^{(n+1)} = \langle \Psi | \{ W (R^{\odot}W)^n \}_c | \Psi \rangle.$$

Let us analyse the significance of these statements for the size extensivity of the calculated energies, understood as the proper dependence of the energy on the size of the system in the limit of non-interacting fragments.

Let us analyse what happens with the <sup>general</sup> connected quantity of the  $\langle \Psi | \{ W (R^{\odot}W)^n \}_c | \Psi \rangle$  or  $\{ (R^{\odot}W)^n \}_c |\Psi\rangle$  type

(let us call this quantity ~~connected~~  $A$ ), when a given system separates into non-interacting fragments

$A, B, \dots$

$$\text{System} \rightarrow A + B + \dots = \sum_{\text{C}} \text{C}_{\text{Segments}}$$

-617-

First of all, in the limit of non-interacting fragments, the spin-orbitals of the entire system become the spin-orbitals of subsystems,

$$|\psi\rangle \Rightarrow |\psi_A\rangle, |\psi_B\rangle, \dots (|\psi_C\rangle \text{ in general}),$$

or  $|\psi_{\text{PA}}\rangle, |\psi_{\text{PB}}\rangle, \dots$  (this, of course,

depends on how we calculate the spin-orbitals, but with the judicious choice of spin-orbitals, using, say ~~poorly~~ <sup>poorly</sup> estimated Hartree-Fock spin-orbitals, we can guarantee that the spin-orbitals of a system of noninteracting fragments are spin-orbitals of subsystems)

Let me illustrate this statement by employing the (unrestricted) Hartree-Fock equations. We can easily show that the Hartree-Fock spin-orbitals of noninteracting fragments (let us focus on molecular fragments), satisfying

$$\left[ -\frac{1}{2} \Delta_i + \sum_{j \in C} \frac{\zeta_j}{r_{ij}} \right] \psi_{ic}(x_i) = \text{coordinates of electron } i$$

electronic coordinate  
 nuclei of C      distance between electron i and nuclear of  
 $\int \psi_{jc}^*(x_2) \psi_{jc}(x_2) dx_2 \quad \psi_{ic}(x_i)$

$$+ \sum_{j \in C \text{ core. in }} \int \psi_{jc}^*(x_2) \psi_{jc}(x_2) dx_2 \quad \psi_{ic}(x_i)$$

68-

$$- \sum_{\substack{j \in E_{\text{occ,in}} \\ C}} \int \frac{\chi_{jC}^*(x) \chi_{iC}(x)}{r_{12}} dx_2 \chi_{jC}(x_1)$$

$$= \varepsilon_{iC} \chi_{iC}(x_1),$$

for each subsystem  $C$  ( $C=A, B, \dots$ ),

satisfies the <sup>H-F</sup> equations for the whole system,

~~$$-\frac{1}{2} \Delta_i = \sum_D \sum_{\substack{j \in E_{\text{occ,in}} \\ C}} \frac{\chi_{jC}(x_1)}{r_{12}} \int \chi_{jC}^*(x_2) \chi_{jC}(x_2) dx_2 \chi_{iC}(x_1)$$~~


$$(1) \quad -\frac{1}{2} \Delta_i = \sum_D \sum_{\substack{j \in E_{\text{occ,in}} \\ C}} \frac{\chi_{jC}(x_1)}{r_{12}} \int \chi_{jC}^*(x_2) \chi_{iC}(x_1)$$

summation  
over all subsystems      nuclei of subsystem D

$$(2) + \sum_D \sum_{\substack{j \in E_{\text{occ,in}} \\ D}} \int \frac{\chi_{jD}^*(x_2) \chi_{jD}(x_2)}{r_{12}} dx_2 \chi_{iC}(x_1)$$

$$(3) + \sum_D \sum_{\substack{j \in E_{\text{occ,in}} \\ D}} \int \frac{\chi_{jD}^*(x_2) \chi_{iC}(x_2)}{r_{12}} dx_2 \chi_{jD}(x_1)$$

$\gamma_{ic} \gamma_{ic}(x_i)$ , when distances between subsystems,  $R_{CD} \rightarrow \infty$ .



$\gamma_{ij}$  - S-nuclei  
on C, D

$R_C, R_D$  - vectors  
of coordinates  
of centers of  
C and D

(1):  ~~$\gamma_{ic}$  is localized on C so that to give a non zero value~~  
 ~~$r_1 \approx R_C$~~  For  $D \neq C$ ,  
 ~~$r_{15} \rightarrow \infty$~~

(1) :  $\gamma_{ic}$  is localized on C, so that  
 $r_1 \approx R_C$  to give a non-zero value of  
 $\gamma_{ic}(x_i)$ . For  $D \neq C$ ,  $r_{15} \rightarrow \infty$   
 if  $R_{CD} \rightarrow \infty$  and  $r_1 \approx R_C$ , so that

-620-

(1) ~~reduces~~ reduces to

$$\left[ -\frac{1}{2} \Delta_1 - \sum_{j \in C} \frac{\chi_j}{r_{ij}} \right] \chi_{iC}(x)$$

(2):  $\chi_{jD}$  is localized on D and  $\chi_{iC}$  is  
localized on C. Thus,  $r_j \approx R_C$  and

$r_j \approx R_D$  to give nonzero values of  $\chi_{iC}$   
and  $\chi_{jD}$ . For  $C \neq D$ ,  $R_D \rightarrow \infty$ ,

$r_{12} \rightarrow \infty$ , so that (2) ~~reduces~~ reduces to

$$\sum_{j \in C} \int \frac{\chi_{jC}(x_2)^* \chi_{jC}(x_2)}{r_{12}} dx_2 \chi_{iC}(x).$$

(3): ~~Again~~ In this case, if  $D \neq C$ ,

$$\chi_{jD}(x_2) \chi_{iC}(x_2) \rightarrow 0 \quad \text{if}$$

$R_D \rightarrow \infty$ . Thus, (3) ~~reduces~~

reduces to  $\sum_{j \in C \text{ or } iC} \int \frac{\chi_{jC}^*(x_2) \chi_{iC}(x_2)}{r_{12} + \chi_{jC}(x_2)} dx_2$

-621-

In other words, the H-F equations for the entire system reduce to


$$\begin{aligned}
 & \left[ -\frac{1}{2} \Delta_1 - \sum_{j \in C} \frac{\gamma_j}{\gamma_{ij}} \right] \chi_{ic}(x_1) \\
 & + \sum_{j \in C} \frac{\int \chi_{jc}^*(x_2) \chi_{jc}(x_2) dx_2}{\gamma_{12}} \chi_{ic}(x_1) \\
 & - \sum_{j \in C} \frac{\int \chi_{jc}^*(x_2) \chi_{jc}(x_2) dx_2}{\gamma_{12}} \chi_{jc}(x_1) \\
 & = \varepsilon_{ic} \chi_{ic}(x_1),
 \end{aligned}$$

i.e. to the equations for individual subsystems.

Thus, we can indeed assume that the spin-orbits of a system of noninteracting fragments are spin-orbits of these fragments.

Now, let us return to quantity 1.

This quantity consists of the connected hexagons, schematically, the



diagrams obtained by connecting some members of  $W$  vertices.  $P_C, q_D, V_E, S_F$  are the symbobols of subsystems  $C, D, E, F$ . Algebraic expressions will contain matrix elements

$$\langle P_C q_D | \tilde{v} | V_E S_F \rangle \xrightarrow[\text{assume } D = \frac{1}{r_{12}}]{\Rightarrow} = \int \psi_{P_C}(x_1)^* \psi_{q_D}(x_2)^* \psi_{V_E}(x_1) \psi_{S_F}(x_2) dx_2$$

First of all, in the noninteracting limit,

$$\psi_{P_C}(x)^* \psi_{V_E}(x_1) \text{ and}$$

-623-

$$\gamma_{qD}(x_2)^* \gamma_{SF}(x_2)$$

removal of  $C \neq E$  and  $D \neq F$ , (because of  
a local character of symbols).

Thus,  $C = E$ ,  $D = F$ , and we are left  
with terms such as

$$\int_{x_1} \underbrace{\gamma_{PC}(x_1)^* \gamma_{qD}(x_2)^* \gamma_{rc}(x_1) \gamma_{SD}(x_2)}_{dx_1 dx_2}$$

Now, if  $C \neq D$  and  $R_D \rightarrow \infty$ ,

we have:

$$\gamma_1 \approx R_C, \gamma_2 \approx R_D,$$

to give momenta  $\gamma_{PC}(x_1)$  and  $\gamma_{qD}(x_2)$

(or  $\gamma_{rc}(x_1)$  and  $\gamma_{SD}(x_2)$ ), in which case


$\gamma_2 \rightarrow \infty$  and the integral vanishes. Thus,

we can only have integrals with  $C = D = E = F$ ,

$$\langle P_C q_C | \bar{v} | r_C s_C \rangle,$$

and, in diagrams,

$$\sum_{c \in C_r} \sum_{p_c q_c r_c s_c}$$



which means that

$$\Delta = \sum_C \Delta_C, \text{ where } \Delta_C$$

is a quantity  $\Delta$  written for fragment  $C$ .

For the connected quantity, we have

$$\Delta = \sum_C \Delta_C, \text{ where}$$

$\Delta_C$  is  $\Delta$  written (or drawn) for fragment  $C$ .

Clearly, since path-variable of

different fragments satisfy the ZDO condition,

$$\chi_{pc}(x) * \chi_{qd}(x)$$

is zero for  $C \neq D$ ,

-625-

No.

They are also orthogonal, and excited configurations for different fragments are orthogonal, too.

Thus,

$$\langle \psi_C | \psi_D \rangle = \delta_{CD} \quad \text{for } C \neq D.$$

In particular,

$$k_S^{(n)} = \langle \psi_0 | \hat{H} (R^{(0)})^n \} \psi_0 \rangle$$

$$= k_0^{(n)}(A) + k_0^{(n)}(B) + \dots,$$

where

$$k_0^{(n)}(A) = \langle \psi_0^{(A)} | \{ \hat{H}^{(A)} (R^{(0),A})^n \} | \psi_0^{(A)} \rangle$$

$$k_0^{(n)}(B) = \langle \psi_0^{(B)} | \{ \hat{H}^{(B)} (R^{(0),B})^n \} | \psi_0^{(B)} \rangle$$

etc.,

are energy corrections for the fragments.

(MBPT)

FINITE-ORDER CALCULATIONS LEAD TO  
SIZE EXTENSIVE RESULTS FOR ENERGIES  
(but not for wave functions!).

Connected cluster theorem  
coupled-cluster ansatz for  $\langle R \rangle$  were mentioned


We know that

$$\langle R^{(n)} \rangle = \{(R^{(n)})^n\}, \langle \mathbb{1} \rangle,$$


These L denotes all linked diagrams ~~with~~,  
including EPV terms. We can classify all  
linked terms according to the number of connected  
components in a diagram:

$L_r$  - all linked diagrams with  $r$   
connected components.


Examples:



are the  $r=1$  diagrams (class  $L_1$ )



are the  $r=2$  diagrams (class  $L_2$ ),



, etc. are the  $r=3$  L-diags (class  $L_3$ ).

-627-

Thus, we can write

$$|\tilde{\Psi}_0^{(n)}\rangle = \sum_{r=1}^n \{(\tilde{R}^0 W)^n\}_{L_r} |\tilde{\Psi}\rangle,$$

bracketed diagrams with  
 r connected components

and  $|\tilde{\Psi}_0\rangle = \sum_{n=0}^{\infty} |\tilde{\Psi}_0^{(n)}\rangle$

$$= |\tilde{\Psi}\rangle + \sum_{n=1}^{\infty} \sum_{r=1}^n \{(\tilde{R}^0 W)^n\}_{L_r} |\tilde{\Psi}\rangle$$

Among classes  $L_r$ , we have class  $L_1$  of  
the connected diagrams. Let us define the  
**CLUSTER OPERATOR  $\hat{T}$**  as the sum of  
all connected ( $L_1$ ) components of  $|\tilde{\Psi}_0\rangle$ .

$$\begin{aligned} \hat{T}|\tilde{\Psi}\rangle &= \sum_{n=1}^{\infty} \{(\tilde{R}^0 W)^n\}_{L_1} |\tilde{\Psi}\rangle \\ &= \sum_{n=1}^{\infty} \{(\tilde{R}^0 W)^n\}_C |\tilde{\Psi}\rangle \end{aligned}$$


-628-

Clearly, connected diagrams  $\{(R^{(0)}H)^n\}_C | \bar{\phi} \rangle$  have certain number of external lines extending to the left ( $2k$  lines,  $k=1, 2, \dots, N$  for an  $N$ -electron system), thus, we can write

$$T|\bar{\phi}\rangle = \sum_{n=1}^{\infty} \sum_{k=1}^N \{(R^{(0)}H)^n\}_{C,k} | \bar{\phi} \rangle,$$

where  $2k$  is the number of external lines in diagrams in  $\{(R^{(0)}H)^n\}_{C,k} | \bar{\phi} \rangle$  (it may happen that for some  $n$  values, not all  $k$  values are possible, in which case  $\{(R^{(0)}H)^n\}_{C,k}$  is a zero form).

Clearly,  $k$  is the excitation number; a diagram



is a linear combination of

$$E_{i_1 \dots i_k}^{a_1 \dots a_k} = X_{a_1}^+ X_{i_1}^- - X_{a_k}^+ X_{i_k}^-$$

-62-

or  $\langle \bar{\phi}_{i_1 \dots i_k}^{d_1 \dots d_n} \rangle = \langle \bar{\phi}_{i_1 \dots i_k}^{d_1 \dots d_n} | \bar{\phi}_0 \rangle$ .

Thus,

$$\begin{aligned} T|\bar{\phi}_0\rangle &= \sum_{k=1}^N \sum_{n=1}^{\infty} \{(R^{(0)H})^n\}_{C_k} |\bar{\phi}_0\rangle \\ &= \sum_{k=1}^N T_k |\bar{\phi}_0\rangle, \end{aligned}$$

or  $T = \sum_{k=1}^N T_k$ , where

$$T_k = \sum_{n=1}^{\infty} T_k^{(n)}, \quad \forall n$$

$$T_k^{(n)}|\bar{\phi}_0\rangle = \{(R^{(0)H})^n\}_{C_k} |\bar{\phi}_0\rangle.$$

all connected diagrams  
resulting from n vertices with  
2k external lines

$T_k$  is a  $k$ -body cluster component,

$T_k^{(n)}$  is the  $n$ -order contribution to  $T_k$ .

-630

(15)

Clearly,

some coefficients resulting from  $\{(\mathbf{x}_k^T \mathbf{W})^m\}$

$$T_k^{(n)}(\mathbf{x}_{\alpha_0}) = \frac{1}{k!} \sum_{\substack{\alpha_1 \dots \alpha_k \\ i_1 \dots i_k}} \langle \alpha_1 \dots \alpha_k | t_k^{(n)} | i_1 \dots i_k \rangle \\ \times \underbrace{\sum_{i_1 \dots i_k}^{\alpha_1 \dots \alpha_k}}_{N[\mathbf{x}_{\alpha_1}^T \mathbf{x}_{i_1} - \mathbf{x}_{\alpha_k}^T \mathbf{x}_{i_k}]}.$$

$$= \left(\frac{1}{k!}\right)^2 \sum_{\substack{\alpha_1 \dots \alpha_k \\ i_1 \dots i_k}} \langle \alpha_1 \dots \alpha_k | t_k^{(n)} | i_1 \dots i_k \rangle_A \\ \times N[\mathbf{x}_{\alpha_1}^T \mathbf{x}_{i_1} - \mathbf{x}_{\alpha_k}^T \mathbf{x}_{i_k}],$$

$$T_k = \frac{1}{k!} \sum_{\substack{\alpha_1 \dots \alpha_k \\ i_1 \dots i_k}} \langle \alpha_1 \dots \alpha_k | t_k | i_1 \dots i_k \rangle \\ \times N[\mathbf{x}_{\alpha_1}^T \mathbf{x}_{i_1} - \mathbf{x}_{\alpha_k}^T \mathbf{x}_{i_k}]$$

$$= \left(\frac{1}{k!}\right)^2 \sum_{\substack{\alpha_1 \dots \alpha_k \\ i_1 \dots i_k}} \langle \alpha_1 \dots \alpha_k | t_k | i_1 \dots i_k \rangle_A \\ \times N[\mathbf{x}_{\alpha_1}^T \mathbf{x}_{i_1} - \mathbf{x}_{\alpha_k}^T \mathbf{x}_{i_k}]$$

then

$$\langle \alpha_1 \dots \alpha_k | t_k | i_1 \dots i_k \rangle = \sum_n \langle \alpha_1 \dots \alpha_k | t_k^{(n)} | i_1 \dots i_k \rangle,$$

and

$$\langle \alpha_1 \dots \alpha_k | t_k | i_1 \dots i_k \rangle_A = \sum_{R \in S_k} \langle \alpha_1 \dots \alpha_k | t_k | i_R \dots i_{R_k} \rangle.$$

-631-

For example (pair-cluster operator or the doubly excited cluster component)

$$\begin{aligned} T_2 &= \frac{1}{4} \sum_{cjob} \langle ab | h_2 | ij \rangle_A N [X_a^+ X_i^- X_b^+ X_j^-] \\ &= \frac{1}{2} \sum_{cjob} \langle ab | h_2 | ij \rangle_A N [X_a^+ X_i^- X_b^+ X_j^-] \underbrace{\varepsilon_{ij}^{ab}}_{\langle ab | ij \rangle} \end{aligned}$$

These

$$\langle ab | h_2 | ij \rangle_A = \langle ab | h_2 | ij \rangle - \langle ab | h_2 | ji \rangle$$

$$T_2 = \sum_{n=1}^{\infty} T_2^{(n)}, \text{ where}$$

$$T_2^{(n)} \underset{\oplus}{\text{Diagram}} = \{ (R^{\otimes n} H)^n \}_{C_2} |\bar{E}\rangle$$

connected diagrams with  
4 external lines.

In the lowest order,

$$T_2^{(1)} \underset{\oplus}{\text{Diagram}} = \{ (R^{\otimes 1} H)^1 \}_{C_2} |\bar{E}\rangle$$

$$= \cancel{\text{Diagram}} = \frac{1}{4} \sum_{cjob} \frac{\langle ab | ij \rangle_A}{\varepsilon_i - \varepsilon_a + \varepsilon_j - \varepsilon_b} \times \cancel{\langle ab | ij \rangle}$$

$$= \sum_{\substack{i,j \\ a,b}} \frac{\langle a b | \delta | i j \rangle_A}{\varepsilon_i - \varepsilon_a + \varepsilon_j - \varepsilon_b} \begin{cases} | \overset{\text{ab}}{\underset{\text{ij}}{\longleftrightarrow}} \rangle \\ \langle \overset{\text{ab}}{\underset{\text{ij}}{\longleftrightarrow}} | b \rangle \end{cases}$$

On the other hand

$$T_2^{(0)} | \Phi_0 \rangle = \frac{1}{4} \sum_{a,b} T_{ab} | t_2^{(0)} | ij \rangle_A E_{ij}^{ab} | \Phi_0 \rangle$$

$$= \sum_{\substack{i,j \\ a,b}} T_{ab} | t_2^{(0)} | ij \rangle_A | \overset{\text{ab}}{\underset{\text{ij}}{\longleftrightarrow}} \rangle$$

so that

$$\boxed{\langle a b | t_2^{(0)} | ij \rangle_A = \frac{\langle a b | \delta | i j \rangle_A}{\varepsilon_i - \varepsilon_a + \varepsilon_j - \varepsilon_b}}$$

Another example:

$T_1$  (the Hartree-Fock case):

$$T_1 = \sum_{\varepsilon_i, a} \langle a | t_1 | i \rangle E_i^a ,$$

$$T_1 = \sum_{n=1}^{\infty} T_1^{(n)}, \quad \text{where}$$

$$T_1^{(n)} | \Phi \rangle = \underbrace{\{(R^{(n)} W)^n\}_{c_1}}_{\text{connected L-mesh with 2 external lines}} | \Phi \rangle$$

-633-

In the H-F case,  $W = V_N$

$$T_i^{(n)}(\mathbb{E}) = \{(R^{\otimes n} V_N)^n\}_{C_1}(\mathbb{E}).$$


The lowest orders:

$n=1$ :

$$T_i^{(1)}(\mathbb{E}) = \{(R^{\otimes 1} V_N)\}_{C_1}(\mathbb{E}) \\ = O(m \text{ diagrams})$$

$n=2$ :

$$T_i^{(2)}(\mathbb{E}) = \{(R^{\otimes 2} V_N)^2\}_{C_2}(\mathbb{E})$$



$$= -\frac{1}{2} \sum_{a,i,m,n,e} \frac{\langle e_a | \hat{v} | m n \rangle_A \langle m n | \hat{v} | e_i \rangle_A}{(\epsilon_i - \epsilon_a)(\epsilon_m + \epsilon_n - \epsilon_a - \epsilon_e)} \langle \Phi_i^a \rangle$$

$$+ \frac{1}{2} \sum_{a,i,m,ef} \frac{\langle m a | \hat{v} | e f \rangle_A \langle e f | \hat{v} | m i \rangle_A}{(\epsilon_i - \epsilon_a)(\epsilon_m + \epsilon_i - \epsilon_e - \epsilon_f)} \langle \Phi_i^a \rangle$$

$$= \sum_{a,i} \langle \alpha | t_i^{(2)} | i \rangle \langle \Phi_i^a \rangle, \text{ so that}$$

$$\langle \alpha | t_i^{(2)} | i \rangle = \frac{1}{2} \sum_{ef,m} \frac{\langle m a | \hat{v} | e f \rangle_A \langle e f | \hat{v} | m i \rangle_A}{(\epsilon_i - \epsilon_a)(\epsilon_m + \epsilon_n - \epsilon_e - \epsilon_f)}$$

$$- \frac{1}{2} \sum_{mne} \frac{\langle e a | \hat{v} | m n \rangle_A \langle m n | \hat{v} | e i \rangle_A}{(\epsilon_i - \epsilon_a)(\epsilon_m + \epsilon_n - \epsilon_a - \epsilon_e)}$$

As we can see, ~~in~~ in the H-F case,

$$T_1 = T_1^{(2)} + \dots$$


$$T_2 = T_2^{(1)} + \dots,$$

$T_2$  is <sup>a lot</sup> more important than  $T_1$ .

- 635 -

We can use similar analysis to show that

$$T_3 = T_3^{(2)} + \dots$$



$$T_4 = T_4^{(3)} + \dots$$

Now, we will prove a CONNECTED CLUSTER THEOREM,

$$|\Psi\rangle = e^T |\Phi\rangle,$$

where

$$T|\Phi\rangle = \sum_{n=1}^{\infty} \{(R^{(0)H})^n\} |\Phi\rangle.$$

Proof is based on the ~~expansion~~ fact that

$$\sum_{n=r}^{\infty} \{(R^{(0)H})^n\} |\Phi\rangle =$$

$$= \frac{1}{r!} T^r |\Phi\rangle. \quad (A)$$

we must have  
 at least  $r$   
 $H$  vertices  
 to get  $r$   
 diagonals

Let us prove Eq. (A). The  $r=1$  case is

Let us,

$$\begin{aligned} & \sum_{n=1}^{\infty} \{(R^{\otimes} W)^n\}_L |\emptyset\rangle = \\ &= \sum_{n=1}^{\infty} \{(R^{\otimes} W)^n\}_C |\emptyset\rangle \\ &\equiv T|\emptyset\rangle = \frac{1}{1!} T^1 |\emptyset\rangle. \end{aligned}$$


The  $r=2$  case:

$$\begin{aligned} & \sum_{n=2}^{\infty} \{(R^{\otimes} W)^n\}_{L_2} |\emptyset\rangle \\ &= \sum_{n=2}^{\infty} \sum_{[A]} \sum_{t_1 < t_2} \{(R^{\otimes} W)^n\}_{\underbrace{[A]_{t_1} [A]_{t_2}}} |\emptyset\rangle \end{aligned}$$

[diagrams with equivalent connected components]



+ all time versions leading to nonequivalent diagrams (not all time versions, since we can get a given diagram as two time versions; cf. Re)



case, where there are 2 time versions, but only 1 is needed, or the



case, where there are 6 time versions, but only 3 are nonequivalent;

The nonequivalent time versions are produced by having a symbiotic ordering of  $[A]_{t_1}, [A]_{t_2}$  diagrams,  $t_1 < t_2$ , to avoid

-637-

repetitions]


$$+ \sum_{n=2}^{\infty} \sum_{[A]_G [B]_G} \sum_{t_1, t_2} \{ (R^{(0)} W)^n \}_{[A]_{t_1} [B]_{t_2}}$$

[diagrams with different connected components]




+ all time versions.  
(in this case,  $\boxed{A}$  and  $\boxed{B}$  are different, so that we need all time versions to get all diagrams of a given type)

To eliminate repetitions, we "order" the connected components in some way; this is important since




examples:

The



In the above notation,  $[A]_{t_1}, [B]_{t_2}$  is a time version



of a given diagram  $\Sigma$  consisting of components  $[A]$  and  $[B]$ .

-638-

123

We assume that components  $[A]$  and  $[B]$  are connected (the  $\angle \cos$ ). Clearly, both components have some external lines (otherwise, we would get an unlinked contribution).

We obtain,

$$\sum_{n=2}^{\infty} \{ (R^{\odot} H)^n \}_{L_2} | \Phi_0 \rangle$$

$$= \sum_{n=2}^{\infty} \sum_{[A]} \sum_{t_1 < t_2} \{ (R^{\odot} H)^n \}_{[A]_{t_1}, [A]_{t_2}} | \Phi \rangle$$

$$+ \sum_{n=2}^{\infty} \sum_{[A] < [B]} \sum_{t_1 < t_2} \{ (R^{\odot} H)^n \}_{[A]_{t_1}, [B]_{t_2}} | \Phi \rangle$$

$$= \frac{1}{2} \sum_{n=2}^{\infty} \sum_{[A]} \sum_{t_1, t_2} \{ (R^{\odot} H)^n \}_{[A]_{t_1}, [A]_{t_2}} | \Phi \rangle$$

( $t_1$  cannot be  $\leq$  any  $t_2$ )

(  
↑  
we include ALL time versions; to avoid repetitions  
we must include a factor of  $\frac{1}{2}$ , as in

$$\begin{array}{c} \text{Diagram A} \\ = \end{array} \frac{1}{2} \left\{ \begin{array}{c} \text{Diagram B} \\ + \end{array} \right\}$$

-638-

24

$$+ \frac{1}{2} \sum_{n=2}^{\infty} \sum_{[A] \neq [B]} \sum_{t_1, t_2} \{(R^{(Q_W)})^n\}_{[A]_G, [B]_G} | \Psi \rangle$$

$$= \frac{1}{2} \sum_{n=2}^{\infty} \sum_{[A], [B]} \sum_{t_1, t_2} \{(R^{(Q_W)})^n\}_{[A]_G, [B]_G} | \Psi \rangle$$

All time versions of the L diagrams  
consisting of process  $[A]$  and  $[B]$

$$= \frac{1}{2} \sum_{k=1}^{\infty} \sum_{l=1}^{\infty} \sum_{[A], [B]} \sum_{t_1, t_2} \{(R^{(Q_W)})^k (R^{(P_W)})^l\}_{[A]_G, [B]_G} | \Psi \rangle$$

$\uparrow$   $\uparrow$   
k vertices W l vertices W

Factorization

Lemma

$$\frac{1}{2} \sum_{k=1}^{\infty} \sum_{l=1}^{\infty} \sum_{[A]} \sum_{[B]} \{(R^{(Q_W)})^k\}_{[A]} \times$$

$\uparrow$   
connected sl-mes  
[A]

$$\times \{(R^{(P_W)})^l\}_{[B]} | \Psi \rangle$$

connected sl-mes

$$= \frac{1}{2} \sum_{k=1}^{\infty} \sum_{[A]} \{(R^{(Q_W)})^k\}_{[A]} \underbrace{\sum_{l=1}^{\infty} \{(R^{(P_W)})^l\}_{[B]} | \Psi \rangle}_{T}$$

-640-

$$= \frac{1}{2} T^2 |\tilde{\phi}_0\rangle.$$

Thus,  $\sum_{n=2}^{\infty} \{(R^{(0)}H)^n\}_{2,2} |\tilde{\phi}_0\rangle = \frac{1}{2!} T^2 |\tilde{\phi}_0\rangle.$

For a general  $r$  case we just recognize that

$$\sum_{n=r}^{\infty} \{(R^{(0)}H)^n\}_{2,r} |\tilde{\phi}_0\rangle =$$

$$= \frac{1}{r!} \sum_{n=r}^{\infty} \sum_{[A_1], \dots, [A_r]} \sum_{t_1, \dots, t_r} \{(R^{(0)}H)^n\}_{[A_1], \dots, [A_r], t_r} |\tilde{\phi}_0\rangle$$

to eliminate repetitions, e.g.  
in the  $n=2$   
case

connected pieces  $[A_1] \rightarrow [A_r]$

diagram composed of  
 $[A_1], [A_2], \dots, [A_r]$

$$= \frac{1}{r!} \sum_{k_1=1}^{\infty} \dots \sum_{k_r=1}^{\infty} \sum_{[A_1], \dots, [A_r]} \sum_{t_1, \dots, t_r}$$

$$\{(R^{(0)}H)^{k_1}, \dots, (R^{(0)}H)^{k_r}\}_{[A_1], \dots, [A_r], t_r} \xrightarrow{k_i \text{ vertices}} |\tilde{\phi}_0\rangle$$

64(-)

Factorization  
lemma

$$\frac{1}{r!} \sum_{k_1=1}^{\infty} \{(R^{\odot} W)^{k_1}\}_{[A_1]} \times$$

$$\times \dots \times \sum_{k_r=1}^{\infty} \{(R^{\odot} W)^{k_r}\}_{[A_r]} \quad | \langle \bar{s} \rangle$$


connected with ext. lines

$$= \frac{1}{r!} T^r |\bar{s}\rangle.$$

(W<sub>r</sub>)

$$|\bar{s}\rangle = \sum_{n=1}^{\infty} \sum_{r=1}^n \{(R^{\odot} W)^n\}_{[A_r]} |\bar{s}\rangle$$

$$= |\bar{s}\rangle + \sum_{r=1}^{\infty} \sum_{n=r}^{\infty} \{(R^{\odot} W)^n\}_{[A_r]} |\bar{s}\rangle$$



$$= |\bar{\Psi}_0\rangle + \sum_{n=1}^{\infty} -\frac{1}{n!} \text{Tr} [\bar{\Psi}] = e^T [\bar{\Psi}_0],$$

Anch completes the proof.

Other arguments in favor of

$$|\bar{\Psi}_0\rangle = e^T [\bar{\Psi}_0].$$

We know that the exact wave function

$$|\bar{\Psi}\rangle = c_0 |\bar{\Psi}_0\rangle + \sum_{i,a} c_a^i |\bar{\Psi}_i^a\rangle$$

$$+ \sum_{\substack{i < j \\ a < b}} c_{ab}^{ij} |\bar{\Psi}_{ij}^{ab}\rangle + \dots$$

$$= c_0 |\bar{\Psi}_0\rangle + \sum_{r=1}^N \hat{c}_r |\bar{\Psi}_r\rangle,$$

where

$$\hat{c}_r |\bar{\Psi}_r\rangle = \sum_{\substack{i_1 < \dots < i_r \\ a_1 < \dots < a_r}} c_{a_1 \dots a_r}^{i_1 \dots i_r} |\bar{\Psi}_{i_1 \dots i_r}^{a_1 \dots a_r}\rangle.$$

are the  $n$ -body creation operators.

-643-

Clearly,

$$C_{\alpha_1 \dots \alpha_r}^{i_1 \dots i_r} = \langle \Phi_{i_1 \dots i_r}^{\alpha_1 \dots \alpha_r} | \tilde{\Psi} \rangle .$$

Since

$\langle \Phi_{i_1 \dots i_r}^{\alpha_1 \dots \alpha_r} \rangle$  are antisymmetric with respect to permutations of  $i_1, \dots, i_r$  or  $\alpha_1, \dots, \alpha_r$ , the same property applies to  $C_{\alpha_1 \dots \alpha_r}^{i_1 \dots i_r}$ . (In particular, if two  $i$ 's or two  $\alpha$ 's are identical,  $C_{\alpha_1 \dots \alpha_r}^{i_1 \dots i_r} = 0$ ), so that

$$\hat{C}_r |\tilde{\Psi}\rangle = \left(\frac{1}{r!}\right)^2 \sum_{\substack{i_1 \dots i_r \\ \alpha_1 \dots \alpha_r}} C_{\alpha_1 \dots \alpha_r}^{i_1 \dots i_r} \langle \Phi_{i_1 \dots i_r}^{\alpha_1 \dots \alpha_r} \rangle$$

or

$$\hat{C}_r = \left(\frac{1}{r!}\right)^2 \sum_{\substack{i_1 \dots i_r \\ \alpha_1 \dots \alpha_r}} C_{\alpha_1 \dots \alpha_r}^{i_1 \dots i_r} \sum_{\substack{i_1 \dots i_r \\ \alpha_1 \dots \alpha_r}} \langle \Phi_{i_1 \dots i_r}^{\alpha_1 \dots \alpha_r} | \tilde{\Psi} \rangle$$

$$\langle \alpha_1 \dots \alpha_r | \tilde{\Psi} \rangle$$

We can always renormalize  $|\tilde{\Psi}\rangle$  to satisfy

$$\langle \tilde{\Psi} | \tilde{\Psi} \rangle = 1 . \text{ In this}$$

$$\text{case, } S_0 = 1.$$

-644-

Thus,

$$|\Psi\rangle = |\Phi\rangle + \sum_{r=1}^N \hat{c}_r |\Phi\rangle \\ = (I + \hat{C}) |\Phi\rangle$$

Thus

$$\hat{C} = \sum_{r=1}^N \hat{c}_r = \hat{G} + \hat{G}_{+-} + \hat{G}_{NN}$$

The excitation operator.

Let us define,

$$T = \ln(I + \hat{C}) =$$

$$= \sum_{m=1}^{\infty} \frac{(-\hat{C})^{m+1}}{m}$$

In general, operator  $T$  may not exist.

However, in our case,

$$\hat{C}^m = 0 \text{ for } m > N. \\ (\hat{C} \text{ is nilpotent})$$

Thus,

$$T = \sum_{m=1}^{\infty} (-D)^{m+1} \frac{C^m}{m}$$

$$= \sum_{m=1}^{N} (-D)^{m+1} \frac{C^m}{m}$$

$\therefore D$  a self-defined operator, represented by  
a finite expression.

Clearly,  $T$  is an excitation operator,

$$T = C - \frac{C^2}{2} + \frac{C^3}{3} - \dots$$

$$= C_1 + C_2 + C_3 + \dots \neq \frac{(C_1 + C_2 + \dots)^2}{2}$$

$$\Rightarrow T_1 = C_1$$

$$T_2 = C_2 - \frac{C_1^2}{2}, \text{ etc.}$$

$$\text{Since } T = \ln(I+C) \Rightarrow$$

$$I+C = e^T \Rightarrow |n\rangle = e^{T(\mathbb{D})} |n\rangle$$

This does not tell us what is  $T$  from the MBPT point of view, but certainly  $T$  exists.

The advantage of the connected cluster theorem is that  $T$  is defined by a self-defined class of diagrams (connected diagrams) which gives us a deep insight into the structure of a many-electron wave function.

The

$e^T \langle \psi \rangle$  ansatz for  $\langle \psi \rangle$

is the basis of the COUPLED-CLUSTER theory, which is based on ~~solving~~ solving the Schrödinger eqn. for  $T$ , treating  $T$  as an unknown. In practice, we determine  $T$  at a given excitation level,

say  $T = T_1 + T_2$ , and we can plug for  $\langle \alpha(t_i) | i \rangle = t_i^i$  ~~ansatz~~,  $\langle ab | t_2 | ij \rangle = \varepsilon_{ab} t_{ij}$ . cluster amplitudes defining  $T_1, T_2, \text{etc.}$

This has an advantage, since CC <sup>consists</sup> guarantees the correct description of separability of a system into subsystems:



$$H_{AB} = H_A + H_B$$

$$|\Psi_0^{(AB)}\rangle = |\Psi_0^{(A)}\rangle |\Psi_0^{(B)}\rangle$$

(ie we are assembling that reference separates OK).

$$|\Psi_0^{(AB)}\rangle = e^{T^{(AB)}} |\Psi_0^{(AB)}\rangle.$$

$T^{(AB)}$  is connected, so that (cf. Hecker discussion)

$$T^{(AB)} = T^{(A)} + T^{(B)},$$

$$[T^{(A)}, T^{(B)}] = 0.$$

In math,

$$e^{X+Y} = e^X e^Y \text{ if } [X, Y] = 0.$$

Thus,

$$|\Psi^{(AB)}\rangle = e^{T^{(A)} + T^{(B)}} |\Psi_0^{(A)}\rangle |\Psi_0^{(B)}\rangle$$

$$= e^{T^{(A)}} |\Psi_0^{(A)}\rangle e^{T^{(B)}} |\Psi_0^{(B)}\rangle$$

$$= |\Psi^{(A)}\rangle |\Psi^{(B)}\rangle, \text{ which is}$$

a desirable behavior.

The energy,

$$E^{(AB)} = \langle \Psi_0^{(AB)} | H_{AB} e^{T^{(AB)}} |\Psi_0^{(AB)}\rangle$$

$$= \langle \Psi_0^{(A)} | \langle \Psi_0^{(B)} | (H_A + H_B) e^{T^{(A)}} |\Psi_0^{(A)}\rangle e^{T^{(B)}} |\Psi_0^{(B)}\rangle$$

$$= \langle \Psi_0^{(A)} | H_A |\Psi_0^{(A)}\rangle + \langle \Psi_0^{(B)} | H_B |\Psi_0^{(B)}\rangle$$

$$\langle \Psi_0^{(A)} | \Psi_0^{(A)} \rangle = E^{(A)} + E^{(B)}, \text{ which is perfect.}$$

The CC engine generates the correct separability  
and size extensivity of the results.